

COMBINING DATASETS TO OVERCOME SELECTION CAUSED BY CENSORING AND TRUNCATION IN MOMENT BASED MODELS

PAUL J. DEVEREUX AND GAUTAM TRIPATHI

ABSTRACT. Economists and other social scientists often encounter data generating mechanisms (dgm's) that produce censored or truncated observations. These dgm's induce a probability distribution on the realized observations that differs from the underlying distribution for which inference is to be made. If this dichotomy between the target and realized populations is not taken into account, statistical inference can be severely biased. In this paper, we show how to do efficient semiparametric inference in moment condition models by supplementing the incomplete observations with some additional data that is not subject to censoring or truncation. These additional observations, or refreshment samples as they are sometimes called, can often be obtained by creatively matching existing datasets. To illustrate our results in an empirical setting, we show how to estimate the effect of changes in compulsory schooling laws on age at first marriage, a variable that is censored for younger individuals. We also demonstrate how refreshment samples for this application can be created by matching cohort information across census datasets.

1. INTRODUCTION

In applied research, economists and other social scientists often encounter data generating mechanisms (dgm's) that produce censored or truncated observations. These dgm's induce a probability distribution on the resulting data, the *realized population*, that differs from the underlying distribution for which inference is to be made, the *target population*. If this dichotomy between the target and realized populations is not taken into account, statistical inference can be severely biased.

To formulate this mathematically, let the triple (Z^*, f^*, μ^*) describe the target population, where Z^* is a random vector (following usual mathematical convention, “vector” means a column vector) in \mathbb{R}^d that denotes an observation from the target population, and f^* is the unknown density of Z^* with respect to some dominating measure $\mu^* = \otimes_{i=1}^d \mu_i^*$. Note that since Z^* is allowed to have discrete components, the μ_i^* 's need not all be Lebesgue measures. Similarly, let (Z, f, μ) represent the realized population, where Z denotes an observation from the realized population and f its density with respect to a dominating measure $\mu = \otimes_{i=1}^d \mu_i$. The basic inferential problem can now be succinctly stated as: *The model comes from f^* but the data comes from f .*

Date: Preliminary version: October 22, 2004. Comments welcome.

Key words and phrases. Censoring, Empirical likelihood, GMM, Refreshment samples, Truncation.

An earlier version of this paper was circulated under the title “GMM and empirical likelihood with incomplete data”. We thank Yuichi Kitamura and Arthur Lewbel for helpful suggestions. The first author gratefully acknowledges financial support from the NSF.

In this paper, we investigate models where f^* is different from f because some or all coordinates of Z^* are censored, or, truncated. The main objective is to develop new techniques that will allow economists to do efficient semiparametric inference by supplementing the original incomplete observations (the *master* sample) with some additional data (the *refreshment* sample) that are complete, i.e., not subject to the mechanism that created the censored or truncated observations¹. Note that the existence of refreshment samples should not be regarded as being an overly restrictive requirement. As illustrated in Section 6, they can often be constructed by creatively *matching* existing datasets.

The idea of using supplementary or refreshment samples to ensure identification in incomplete, biased sampling, or measurement error models has of course been recognized earlier; see, e.g., Manski and Lerman (1977), Titterton (1983), Titterton and Mill (1983), Vardi (1985), Ridder (1992), Carroll, Ruppert, and Stefanski (1995), Wansbeek and Meijer (2000), Hirano, Imbens, Ridder, and Rubin (2001), Chen, Hong, and Tamer (2004), Tripathi (2003, 2004), and the references therein. However, the use of matching to facilitate moment based inference in overidentified models with incomplete data seems to be new to the literature and the results obtained in this paper cannot be found in any of the references cited here.

Our inferential approach is based on the generalized method of moments (GMM) proposed by Hansen (1982) and empirical likelihood (EL) proposed by Owen (1988). We focus on GMM because its unifying approach and wide applicability has made it the method of choice for estimating and testing nonlinear economic models. Its availability in canned software packages has also added to its popularity with applied economists. An excellent exposition on GMM can be found in Newey and McFadden (1994). We also look at EL because it has lately begun to emerge as a serious contender to GMM; see, e.g., Qin and Lawless (1994), Imbens (1997), Kitamura (1997), Smith (1997), Owen (2001), and the 2002 special issue of the Journal of Business and Economics Statistics.

This paper illustrates the advantages of using supplementary or refreshment samples to handle incomplete data: Many partially observed or missing data models in applied work are identified by assuming that the selection probabilities (i.e., the probability that an observation is fully observed) — or propensity scores, as they are often called — are completely known or can be parametrically modelled; see, e.g., Robins, Rotnitzky, and Zhao (1994), Robins, Hsieh, and Newey (1995), Nan, Emond, and Wellner (2002), and the references therein. However, as noted by Horowitz and Manski (1998), these assumptions lead to stochastic restrictions that are usually not testable. In contrast, since we rely upon a refreshment sample to provide identifying power, the selection probabilities in this paper are fully nonparametric (i.e., completely unrestricted) and, hence, we can avoid the above mentioned non-testability problems. More generally, in this paper we demonstrate how a refreshment sample allows standard GMM and EL based inference, including tests of overidentifying restrictions, with censored or truncated data to go through without imposing parametric, independence, symmetry, quantile, or special regressor restrictions.

¹Refreshment samples for most applications will be much smaller than the master sample. Therefore, for efficiency considerations we combine the two samples rather than use the refreshment sample alone.

Furthermore, there is no need to worry about issues relating to heteroscedasticity (a major concern for parametric models) because GMM and EL automatically produce correct standard errors, and, unlike quantile restriction models, there is no need to restrict attention to applications where only scalar-valued continuously distributed random variables are censored or truncated, or use any nonparametric smoothing procedures to estimate asymptotic variances. Extension to the case where more than one random variable (discrete or continuous) is censored or truncated is straightforward and the usual analogy principle that delivers standard errors for GMM or EL works here as well.

The treatment proposed here is general enough to handle censoring and truncation of some or all coordinates of both endogenous and exogenous variables and the results obtained here are applicable to a large class of potentially overidentified models which nest linear regression as a special case; e.g., the ability to handle IV models permits semiparametric inference in Box-Cox type transformation models using censored or truncated data without imposing parametric or quantile restrictions.

The rest of the paper is organized as follows. In Section 2 we set up the moment based model and describe how to deal with the censoring or truncation of random vectors in a multivariate framework, several examples of which are given throughout the paper (including Section 3). Section 4 shows how censored data can be combined with a refreshment sample to do efficient moment based inference and Section 5 does the same with truncated data. Section 6 contains an interesting application where refreshment samples are obtained by matching census datasets. Section 7 concludes by addressing some topics for future research. Proofs, tables, and figures are all in the appendices.

2. SETUP

The econometric models we consider can be expressed in terms of moment conditions as follows. Let Θ be a subset of \mathbb{R}^p such that

$$H_0 : \mathbb{E}_{f^*}\{g(Z^*, \theta^*)\} = 0 \quad \text{for some } \theta^* \in \Theta, \quad (2.1)$$

where g is a $q \times 1$ vector of known functions with $q \geq p$ and \mathbb{E}_{f^*} denotes expectation with respect to the pdf f^* . Models specified via moment conditions are particularly important for structural estimation: Since economic theory attributes outcomes at the micro level to optimizing behavior on the part of firms or individuals, moment based models arise naturally in microeconometrics as solutions to the first order conditions of the stochastic optimization problems economic agents are assumed to solve.

Well-known statistical examples of (2.1) include linear regression models $Y^* = X^{*\prime}\theta^* + \varepsilon$, where the error term is uncorrelated with the regressors; i.e., $\mathbb{E}_{f^*}\{X^*\varepsilon\} = 0$. Here $g(Z^*, \theta^*) = X^*\{Y^* - X^{*\prime}\theta^*\}$ and $Z^* = (Y^*, X^*)$, where Y^* denotes the endogenous variable and X^* the vector of explanatory variables. This can be extended to nonlinear regression models of the form $Y^* = \psi(X^*, \theta^*) + \varepsilon$, where ψ is known up to θ^* and $\mathbb{E}_{f^*}\{\frac{\partial\psi(X^*, \theta^*)}{\partial\theta}\varepsilon\} = 0$. Other generalizations include linear and nonlinear regression models with endogenous regressors and multivariate simultaneous equations models where the error terms are uncorrelated with some or all of the explanatory variables.

The class of models defined in (2.1) also contains instrumental variables (IV) models. Suppose we have the conditional mean restriction $\mathbb{E}_{Y^*|X^*}\{\tilde{g}(Y^*, X^*, \theta^*)|X^*\} = 0$ w.p.1, where \tilde{g} is a $k \times 1$ vector of known functions and Y^* the vector of endogenous variables. Letting $A(X^*)$ denote a $q \times k$ matrix of instruments, this yields unconditional moment restrictions of the form $\mathbb{E}_{f^*}\{g(Z^*, \theta^*)\} = 0$, where $g(Z^*, \theta^*) = A(X^*)\tilde{g}(Y^*, X^*, \theta^*)$.

If the target variable Z^* is fully observed, then (2.1) is easily handled; see, e.g., Newey and McFadden (1994). But in many cases economists cannot fully observe Z^* . For instance, variables often get censored due to administrative reasons; e.g., government agencies routinely “top-code” income data before releasing it for public use. Similarly, studies investigating the length of unemployment spells can terminate prematurely due to financial constraints before all subjects have found employment.

So suppose that all coordinates of Z^* are right-censored; i.e., instead of observing Z^* we observe the random variable $Z = (Z^{(1)}, \dots, Z^{(d)})_{d \times 1}$, where

$$Z^{(i)} = \begin{cases} Z^{*(i)} & \text{if } Z^{*(i)} < c^{(i)} \\ c^{(i)} & \text{otherwise} \end{cases} \quad \text{for } i = 1, \dots, d$$

and $c = (c^{(1)}, \dots, c^{(d)})$ is a $d \times 1$ vector of known constants².

We allow for the possibility that some components of Z^* may not be censored: If, say, the i th coordinate of Z^* is not subject to the censoring mechanism, simply set $c^{(i)} = \infty$; if the i th and j th coordinates of Z^* , denoted by $Z^{*(i,j)}$, are not subject to censoring, then set $c^{(i,j)} = (\infty, \infty)$; etc.. Hence, in applications where the target variable Z^* can be decomposed into endogenous and exogenous parts as (Y^*, X^*) , we can handle situations where only Y^* is censored (pure endogenous censoring), or only X^* is censored (pure exogenous censoring)³, or only some coordinates of either variables are censored. Left censoring of, say, the i th, j th, and k th coordinates can also be accommodated by replacing $Z^{*(i,j,k)}$ with $-Z^{*(i,j,k)}$ and $c^{(i,j,k)}$ with $-c^{(i,j,k)}$.

Let S^* denote the survival function induced by f^* , i.e., $S^*(\xi) = \Pr_{f^*}(\cap_{i=1}^d \{Z^{*(i)} > \xi^{(i)}\})$, and δ_c the Dirac measure at c , i.e., $\delta_c(A) = \mathbb{I}(c \in A)$, where \mathbb{I} is the indicator function. To keep matters simple, we assume that μ^* does not have an atom at c . This assumption, which can be relaxed at the cost of greater mathematical complexity, is weaker than requiring μ^* to be a Lebesgue measure (the usual assumption made for censored regression models).

If $d = 1$, the density of Z with respect to the dominating measure $\mu = \mu^* + \delta_c$ is given by

$$f(z) = f^*(z)\mathbb{I}(z < c) + S^*(c)\mathbb{I}(z = c). \quad (2.2)$$

²The results obtained in this paper continue to hold in a more general *fixed* censoring framework where the censoring point is modelled as a random variable C with unknown distribution such that C is observed for censored as well as uncensored observations; see, e.g., the application in Section 6.

³The term “exogenous” is, strictly speaking, an abuse of terminology since (2.1) does not involve any conditioning although, as mentioned earlier in the introduction, (2.1) does nest IV models based on conditional moment restrictions. Therefore, the careful reader may want to substitute “censoring (resp. truncation) based on explanatory variables” for “exogenous censoring (resp. truncation)” whenever the latter is encountered.

The density of Z when it is vector valued is also straightforward to derive but requires some additional notation. So let $Z^{*(i,j,k)}$ denote coordinates of Z^* that remain after the i th, j th, and k th ones have been deleted, $f_{-(i,j,k)}^*$ the joint density of $Z^{*(i,j,k)}$, and $S_{i,j,k|-(i,j,k)}^*$ the conditional survival function induced by $f_{i,j,k|-(i,j,k)}^*$, the conditional density of $Z^{*(i,j,k)}|Z^{*(i,j,k)}$. It is then easy to show that for $d > 1$ the density of Z with respect to $\mu = \otimes_{i=1}^d \mu_i$, where $\mu_i = \mu_i^* + \delta_c^{(i)}$, is given by

$$f(z) = f^*(z)\mathbb{I}(z \stackrel{elt}{<} c) + \sum_{r=1}^{d-1} \sum_{i_1=1}^{d-r+1} \sum_{i_2=i_1+1}^{d-r+2} \cdots \sum_{i_r=i_{r-1}+1}^d S_{i_1,\dots,i_r|-(i_1,\dots,i_r)}^*(c^{(i_1,\dots,i_r)}) f_{-(i_1,\dots,i_r)}^*(z^{-(i_1,\dots,i_r)}) \\ \times \mathbb{I}(z^{(i_1,\dots,i_r)} = c^{(i_1,\dots,i_r)}, z^{-(i_1,\dots,i_r)} \stackrel{elt}{<} c^{-(i_1,\dots,i_r)}) + S^*(c)\mathbb{I}(z = c), \quad (2.3)$$

where $\stackrel{elt}{<}$ denotes element-by-element strict inequality; i.e., $\mathbb{I}(z \stackrel{elt}{<} c) = \prod_{i=1}^d \mathbb{I}(z^{(i)} < c^{(i)})$. Of course, $z = c$ still denotes element-by-element equality; i.e., $\mathbb{I}(z = c) = \prod_{i=1}^d \mathbb{I}(z^{(i)} = c^{(i)})$. Note that f has support $(-\infty, c^{(1)}] \times \dots \times (-\infty, c^{(d)}]$ with an atom at c .

Sometimes censoring is so severe that the target variable is completely unobserved outside a certain region. This phenomenon is called truncation; e.g., in many job training programs subjects are allowed entry only if their household income falls below a certain level. If Z^* is a truncated random variable, then instead of observing Z^* we observe

$$Z = \begin{cases} Z^* & \text{if } Z^* \in T \\ \text{unobserved} & \text{otherwise,} \end{cases}$$

where T denotes a known region in \mathbb{R}^d such that Z^* lies in T with positive probability. In this case, the density of Z with respect to μ^* is given by

$$f(z) = \frac{f^*(z)\mathbb{I}(z \in T)}{\int_T f^*(z) d\mu^*}. \quad (2.4)$$

Note that f has support T . As before, we allow for the possibility that some coordinates of Z^* may not be truncated: In typical applications, T will be a rectangle of the form $I_1 \times \dots \times I_d$, where the I_j 's are known fixed intervals. If, say, $Z^{*(i,j,k)}$ are not truncated, then simply let $I_i = I_j = I_k = \mathbb{R}$.

Since in this paper we assume that f^* and, hence, f are completely unknown, censoring or truncation of Z^* creates a fundamental identification problem. To see this, first note that since Z is the observed version of Z^* , the realized density f is identified by definition. However, as is evident from (2.2)–(2.3) and (2.4), the target density $f^*(z)$ cannot be expressed in terms of $f(z)$ for all $z \in \mathbb{R}^d$. In other words, f^* cannot be fully recovered from f ; i.e., f^* is not identified. But if there is no way of going from the realized density (loosely speaking, the “reduced form”) to the target density (the “structural form”), then statistical inference about f^* and, hence, the target cdf $F^*(\xi) = \Pr_{f^*}(Z^* \stackrel{elt}{\leq} \xi)$

is impossible⁴. So the first task is to overcome this lack of identification. Basically, this is done by requiring additional information about f^* .

The information needed to ensure identification of f^* can come in different guises. For instance, the classical solution is to simply assume that f^* is known up to a finite dimensional parameter. Hence, identification of the finite dimensional parameter ensures that f^* is also identified. There is a huge literature on showing how to do maximum likelihood estimation with incomplete observations if a parametric model for f^* is acceptable. See, e.g., Hausman and Wise (1976, 1977), Dempster, Laird, and Rubin (1977), Maddala (1983), Amemiya (1985), and the many references therein. A related methodology was suggested by Heckman (1976, 1979), who used the Gaussian framework to motivate his famous “two step” bias corrected estimator now widely used in applied work. However, as noted in Vella (1998), the maximum likelihood and two step approaches are very sensitive and non-robust to parametric distributional assumptions. Therefore, if it is not feasible to make such strong distributional assumptions, then other approaches that do not require a parametric specification for f^* have to be used instead.

The real challenge in working with censored or truncated data arises when f^* is completely unknown. Most of the literature in this area, see Powell (1994) for a useful survey, seems to have focused on the tobit model with different authors using different assumptions to identify the finite dimensional parameters⁵. For example, Bhattacharya, Chernoff, and Yang (1983), Chamberlain (1986), Duncan (1986), Fernandez (1986), Horowitz (1986, 1988), Cosslett (1987, 1991), Tsui, Jewell, and Wu (1988), Moon (1989), and Honoré and Powell (1994) assume that ε and X^* are independent; Powell (1983, 1984, 1986a), Nawata (1990), Newey and Powell (1990), Buchinsky and Hahn (1998), Chen and Khan (2001), Khan and Powell (2001), and Hong and Tamer (2003) assume knowledge of some quantile of the conditional distribution of ε given X^* ; Powell (1986b), Newey (1988), and Lee (1993a, 1993b) assume that the distribution of $\varepsilon|X^*$ is symmetric. More recently, a different set of assumptions involving the existence of a “special regressor” that is continuously distributed, has large support, and satisfies a certain exclusion restriction, have been used by Khan and Lewbel (2003) to identify slope coefficients in a truncated regression model.

Although independence, symmetry, quantile, or special regressor restrictions suffice to identify finite dimensional parameters, they are not strong enough to permit consistent estimation of the target

⁴Since distribution functions characterize random variables, estimating $F^*(\xi)$ at each $\xi \in \mathbb{R}^d$ determines the probabilistic behavior of Z^* . Efficient estimation of the target cdf is important for bootstrapping from the target population. Brown and Newey (2002) note that when prior information about the target population is available, merely using a consistent estimator of F^* can lead to poor inference from the bootstrap. They recommend that resampling be done using \hat{F}^* , an efficient estimator of F^* that incorporates restrictions imposed by the model (2.1). Estimating the realized cdf under (2.1) is also useful because comparing it with \hat{F}^* can help reveal the extent of bias induced by censoring or truncation. Of course, \hat{F}^* can always be compared with the empirical cdf of the observed data. But since the latter does not take the model into account, it will be less precise (though more robust) than an estimator of the realized cdf under (2.1).

⁵Newey (2001) has some general results regarding the kind of restrictions on the distribution of $\varepsilon|X^*$ that lead to $n^{1/2}$ -consistent estimation of the regression coefficients.

cdf $F^*(\xi)$ at each $\xi \in \mathbb{R}^d$ because they cannot ensure identification of the target density itself. In contrast, we ensure identification of f^* (and, hence, F^*) by assuming that in addition to the original incomplete data (i.e., the master sample) we also have a set of supplementary observations that are unaffected by censoring or truncation (i.e., the refreshment sample).

To be precise, we assume henceforth that the realized random variable Z has density⁶

$$f_e(z) = \begin{cases} K_0 f^*(z) \mathbb{I}(z \stackrel{elt}{\neq} c) + (1 - K_0) f(z) & \text{if } Z^* \text{ is censored} \\ K_0 f^*(z) + (1 - K_0) f(z) & \text{if } Z^* \text{ is truncated,} \end{cases} \quad (2.5)$$

where $K_0 \in (0, 1]$ is a known constant, $\mathbb{I}(z \stackrel{elt}{\neq} c) = \prod_{i=1}^d \mathbb{I}(z^{(i)} \neq c^{(i)})$, and, depending on whether Z^* is censored or truncated, f is given by (2.2)–(2.3) or (2.4), respectively. If Z^* is censored, then f_e is a density with respect to μ and has an atom at c . On the other hand, if Z^* is truncated, then f_e is a density with respect to μ^* .

Loosely speaking, K_0 denotes the probability of sampling from the whole target population without subjecting the observations to censoring or truncation. As a practical example, consider a dataset of 100 observations, 90 of which belong to the original sample of censored observations and the remaining 10 constitute the uncensored supplementary sample. Hence, $K_0 = 0.10$ and this dataset can be regarded as a random sample from f_e . The no censoring or truncation case corresponds to setting $K_0 = 1$. In Sections 4.1 and 5.1 we show how this *enriched* density can be used to identify f^* . Once questions regarding identification are settled, we can proceed with estimating and testing (2.1).

3. EXAMPLES

In this section we look at some examples of censoring and truncation in a multivariate framework. The primary aim is to illustrate what happens in linear models when more than one variable is partially observed. Examples 3.2 and 3.4 are particularly instructive.

Example 3.1 (Censored mean). Suppose we want to estimate $\theta^* = \mathbb{E}_{f^*}\{Z^*\}$, the mean of the target population. Since Z^* is censored from above, instead of a random sample Z_1^*, \dots, Z_n^* from the target density f^* we have a random sample Z_1, \dots, Z_n from the realized density f defined in (2.2) or (2.3). Therefore, the naive estimator $\sum_{j=1}^n Z_j/n$ will not consistently estimate θ^* because $\sum_{j=1}^n Z_j/n \xrightarrow{p} \mathbb{E}_f\{Z\}$ by the weak law of large numbers (WLLN)⁷ but

$$\mathbb{E}_{f^*}\{Z^*\} \neq \mathbb{E}_f\{Z\} = \begin{cases} \mathbb{E}_{f^*}\{Z^* \mathbb{I}(Z^* < c)\} + cS^*(c) & \text{if } d = 1 \\ \mathbb{E}_{f^*}\{Z^* \mathbb{I}(Z^* \stackrel{elt}{<} c)\} + \sum_{r=1}^{d-1} \mathbb{E}_{f^*}\{Z_r^*\} + cS^*(c) & \text{if } d > 1, \end{cases}$$

⁶Since f^* is a density with respect to μ^* , it is only identified up to sets of μ^* -measure zero. Therefore, if Z^* is censored then $f^*(z) \mathbb{I}(z \stackrel{elt}{\neq} c)$ is a μ^* -version of f^* and, hence, $\mathbb{E}_{f^*}\{g(Z^*, \theta^*)\} = 0$ if and only if $\mathbb{E}_{f^*}[g(Z^*, \theta^*) \mathbb{I}(Z^* \stackrel{elt}{\neq} c)] = 0$.

⁷Throughout the paper, all limits are taken as the sample size $n \uparrow \infty$.

where, for any function $h(\cdot)$, the symbol

$$h_r(Z^*) = \sum_{i_1=1}^{d-r+1} \sum_{i_2=i_1+1}^{d-r+2} \dots \sum_{i_r=i_{r-1}+1}^d h(Z^*[i_1, \dots, i_r]) \mathbb{I}(Z^{*(i_1, \dots, i_r)} \stackrel{elt}{>} c^{(i_1, \dots, i_r)}, Z^{*-(i_1, \dots, i_r)} \stackrel{elt}{<} c^{-(i_1, \dots, i_r)})$$

denotes h evaluated at exactly r censored coordinates and $Z^*[i_1, \dots, i_r]$ stands for Z^* with its i_1, \dots, i_r th coordinates replaced by $c^{(i_1)}, \dots, c^{(i_r)}$, respectively, and the remaining coordinates unchanged; i.e., $Z^*[i_1, \dots, i_r] = Z^*|_{Z^{*(i_1, \dots, i_r)} = c^{(i_1, \dots, i_r)}}$. \square

Example 3.2 (Censored linear regression). Let $Y^* = X^{*'}\theta^* + \varepsilon$, where $\mathbb{E}_{f^*}\{X^*\varepsilon\} = 0$. Suppose that both Y^* and X^* are censored. Hence, instead of observing $Z^* = (Y^*, X^*)_{(p+1) \times 1}$ from the target density f^* , we observe $Z = (Y, X)$ from the realized density f defined in (2.3). If we ignore censoring and simply regress Y on X , then θ^* cannot be consistently estimated by the least squares estimator $\tilde{\theta} = (\sum_{j=1}^n X_j X_j')^{-1} \sum_{j=1}^n X_j Y_j$. To see this, observe that the probability limit of $\tilde{\theta}$ is given by

$$\begin{aligned} (\mathbb{E}_f X X')^{-1} (\mathbb{E}_f X Y) &= (\mathbb{E}_{f^*} \{X^* X^{*'} \mathbb{I}(Y^* < c^{(1)}, X^* \stackrel{elt}{<} c^{-(1)}) + \sum_{r=1}^{d-1} (X^* X^{*'})_r + c^{-(1)} c^{-(1)'} S^*(c)\})^{-1} \\ &\quad \times \mathbb{E}_{f^*} \{X^* Y^* \mathbb{I}(Y^* < c^{(1)}, X^* \stackrel{elt}{<} c^{-(1)}) + \sum_{r=1}^{d-1} (X^* Y^*)_r + c^{-(1)} c^{(1)} S^*(c)\}, \quad (3.1) \end{aligned}$$

where $d = p + 1$ and, in the notation introduced in Example 3.1,

$$\begin{aligned} (X^* Y^*)_r &= \sum_{i_1=1}^{d-r+1} \sum_{i_2=i_1+1}^{d-r+2} \dots \sum_{i_r=i_{r-1}+1}^d (X^* Y^*)|_{(Y^*, X^*)^{(i_1, \dots, i_r)} = c^{(i_1, \dots, i_r)}} \\ &\quad \mathbb{I}(Z^{*(i_1, \dots, i_r)} \stackrel{elt}{>} c^{(i_1, \dots, i_r)}, Z^{*-(i_1, \dots, i_r)} \stackrel{elt}{<} c^{-(i_1, \dots, i_r)}), \quad (3.2) \end{aligned}$$

and $(X^* X^{*'})_r$ is obtained by replacing Y^* in (3.2) with $X^{*'}$. Hence, $\text{plim}(\tilde{\theta}) \neq (\mathbb{E}_{f^*} X^* X^{*'})^{-1} (\mathbb{E}_{f^*} X^* Y^*)$.

The special case of pure endogenous censoring, frequently called the tobit or limited dependent variable model in the econometrics literature, is obtained by letting $c^{-(1)} = (\infty, \dots, \infty)$ and using the convention that $0 \cdot \infty = 0$. Doing so, (3.1) implies that

$$\text{plim}(\tilde{\theta}) = \theta^* - \{\mathbb{E}_{f^*} X^* X^{*'}\}^{-1} \mathbb{E}_{f^*} \{X^* (Y^* - c^{(1)}) \mathbb{I}(Y^* > c^{(1)})\} \neq \theta^*,$$

as is well known from tobit theory.

However, a fact that does not seem to be as widely known is that the least squares estimator remains inconsistent even if censoring is purely exogenous. In particular, by letting $c^{(1)} = \infty$ in (3.1),

$$\text{plim}(\tilde{\theta}) = \{\mathbb{E}_{f^*} [X^* X^{*'} \mathbb{I}(X^* \stackrel{elt}{<} c^{-(1)}) + \sum_{r=1}^{d-1} (X^* X^{*'})_r]\}^{-1} \mathbb{E}_{f^*} \{X^* Y^* \mathbb{I}(X^* \stackrel{elt}{<} c^{-(1)}) + \sum_{r=1}^{d-1} (X^* Y^*)_r\} \neq \theta^*,$$

where $(X^*Y^*)_r$ is now equal to

$$\sum_{i_1=2}^{d-r+1} \sum_{i_2=i_1+1}^{d-r+2} \cdots \sum_{i_r=i_{r-1}+1}^d (X^*Y^*) \Big|_{(Y^*, X^*)^{(i_1, \dots, i_r)} = c^{(i_1, \dots, i_r)}} \\ \times \mathbb{I}(X^{*(i_1-1, \dots, i_r-1)} \stackrel{elt}{>} c^{(i_1, \dots, i_r)}, X^{*(i_1-1, \dots, i_r-1)} \stackrel{elt}{<} c^{-(1, i_1, \dots, i_r)}) \quad (3.3)$$

and $(X^*X^*)_r$ follows by replacing Y^* in (3.3) with X^* . Hence, pure exogenous censoring cannot be ignored here. In fact, pure exogenous censoring is not ignorable even if $\mathbb{E}_{f^*}\{X^*\varepsilon\} = 0$ is replaced by the stronger condition $\mathbb{E}_{Y^*|X^*}\{\varepsilon|X^*\} = 0$ w.p.1⁸. However, as shown in Example 3.4, the situation changes if X^* is truncated instead of censored. \square

Example 3.3 (Truncated mean). Again, suppose that we want to estimate the mean of the target population but now Z^* is truncated outside the region T . Since $\mathbb{E}_f\{Z\} = \mathbb{E}_{f^*}[Z^*\mathbb{I}(Z^* \in T)] / \int_T f^*(z) d\mu^*$, as in Example 3.1 the naive estimator is not consistent for $\mathbb{E}_{f^*}\{Z^*\}$. \square

Example 3.4 (Truncated linear regression). Consider the linear model of Example 3.2, but now suppose that instead of being censored, Z^* is truncated outside $T = T_1 \times T_2$. Since the probability limit of the least squares estimator is now given by

$$\text{plim}(\tilde{\theta}) = \{\mathbb{E}_{f^*} X^* X^{*\prime} \mathbb{I}(Y^* \in T_1, X^* \in T_2)\}^{-1} \mathbb{E}_{f^*} \{X^* Y^* \mathbb{I}(Y^* \in T_1, X^* \in T_2)\},$$

it is immediate that $\tilde{\theta}$ is not consistent for θ^* . For pure endogenous truncation, $T_2 = \mathbb{R}^p$. In this case,

$$\text{plim}(\tilde{\theta}) = \{\mathbb{E}_{f^*} X^* X^{*\prime} \mathbb{I}(Y^* \in T_1)\}^{-1} \mathbb{E}_{f^*} \{X^* Y^* \mathbb{I}(Y^* \in T_1)\} \neq \theta^*.$$

Similarly, for pure exogenous truncation, $T_1 = \mathbb{R}$. Hence,

$$\text{plim}(\tilde{\theta}) = \{\mathbb{E}_{f^*} X^* X^{*\prime} \mathbb{I}(X^* \in T_2)\}^{-1} \mathbb{E}_{f^*} \{X^* Y^* \mathbb{I}(X^* \in T_2)\} \neq \theta^*. \quad (3.4)$$

Therefore, even pure exogenous truncation is not ignorable. But unlike Example 3.2, if the identifying assumption $\mathbb{E}_{f^*}\{X^*\varepsilon\} = 0$ is replaced by $\mathbb{E}_{Y^*|X^*}\{\varepsilon|X^*\} = 0$ w.p.1, then from (3.4) it is easy to see that ignoring pure exogenous truncation does not make the least squares estimator inconsistent. \square

4. INFERENCE WITH CENSORED DATA

In this section we show how to do efficient semiparametric inference for moment based models using enriched data (i.e., censored data supplemented with additional observations that are not subject to censoring). We begin by showing that f^* is identified if the realized observations are drawn from the mixture density defined in (2.5).

⁸To get some intuition behind this result, consider the case when $d = 2$; i.e., when $Y^* = X^*\theta^* + \varepsilon$, X^* is scalar, and $c = (\infty, c^{(2)})_{2 \times 1}$. Then, $\text{plim}(\tilde{\theta})$ is equal to

$$\frac{\mathbb{E}_f\{XY\}}{\mathbb{E}_f\{X^2\}} \stackrel{(3.1)}{=} \frac{\mathbb{E}_{f^*}\{X^*Y^*\mathbb{I}(X^* < c^{(2)}) + Y^*c^{(2)}\mathbb{I}(X^* > c^{(2)})\}}{\mathbb{E}_{f^*}\{X^{*2}\mathbb{I}(X^* < c^{(2)}) + [c^{(2)}]^2\mathbb{I}(X^* > c^{(2)})\}} = \frac{\mathbb{E}_{f^*}\{X^{*2}\mathbb{I}(X^* < c^{(2)}) + X^*c^{(2)}\mathbb{I}(X^* > c^{(2)})\}}{\mathbb{E}_{f^*}\{X^{*2}\mathbb{I}(X^* < c^{(2)}) + [c^{(2)}]^2\mathbb{I}(X^* > c^{(2)})\}} \theta^*,$$

where the last equality follows because $\mathbb{E}_{Y^*|X^*}\{Y^*|X^*\} = X^*\theta^*$ w.p.1. Therefore, $\text{plim}(\tilde{\theta}) \neq \theta^*$ and $\tilde{\theta}$ remains inconsistent under pure exogenous censoring even when ε is mean independent of X^* .

4.1. Identification and efficient estimation. By (2.2)–(2.3) and (2.5),

$$f^*(z)\mathbb{I}(z \neq c)^{elt} = \frac{\mathbb{I}(z \neq c)^{elt}}{K_0 + (1 - K_0)\mathbb{I}(z < c)^{elt}} f_e(z). \quad (4.1)$$

Hence, a μ^* -version of f^* is identified since it can be expressed in terms of f_e alone.

Next, since $\mathbb{E}_{f^*}\{g(Z^*, \theta^*)\} = 0$ if and only if $\mathbb{E}_{f_e}[g(Z^*, \theta^*)\mathbb{I}(Z^* \neq c)^{elt}] = 0$, we can use (4.1) to rewrite (2.1) in terms of the enriched density as⁹

$$\mathbb{E}_{f_e}\left\{\frac{g(Z, \theta^*)\mathbb{I}(Z \neq c)^{elt}}{K_0 + (1 - K_0)\mathbb{I}(Z < c)^{elt}}\right\} = 0. \quad (4.2)$$

However, (4.1) also implies that

$$\mathbb{E}_{f_e}\left\{\frac{\mathbb{I}(Z \neq c)^{elt}}{K_0 + (1 - K_0)\mathbb{I}(Z < c)^{elt}}\right\} = 1, \quad (4.3)$$

which represents an additional restriction imposed by the censoring process upon the enriched density. Therefore, an efficient estimator of θ^* based on (4.2) must account for this extra information as well. But since (4.3) can be explicitly solved for K_0 to yield

$$K_0 = \frac{\mathbb{E}_{f_e}\{\mathbb{I}(Z \neq c)^{elt}\mathbb{I}(Z < c)^{elt}\sim\}}{\mathbb{E}_{f_e}\{\mathbb{I}(Z < c)^{elt}\sim\}}, \quad (4.4)$$

where $(Z < c)^{elt}\sim$ denotes the set-complement of the event $(Z < c)^{elt}$ so that $\mathbb{I}(Z < c)^{elt}\sim = 1 - \mathbb{I}(Z < c)^{elt}$, it also reveals that K_0 is an ancillary parameter for estimating θ^* . Therefore, inference regarding θ^* should be done conditional on the estimated value of K_0 (even though K_0 is known!). To do so, we treat K_0 as an unknown parameter and estimate it jointly with θ^* . We explain later how this approach allows us to do conditional inference¹⁰.

⁹It is then easy to show that a sufficient condition for θ^* to be *locally* identified is that the $q \times p$ Jacobian matrix $\mathbb{E}_{f_e}\left\{\frac{\partial g(Z, \theta^*)}{\partial \theta} \frac{\mathbb{I}(z \neq c)^{elt}}{K_0 + (1 - K_0)\mathbb{I}(z < c)^{elt}}\right\}$ exists and is of full column rank. In some models, especially those linear in θ^* , this may be simpler than employing Chamberlain (1986) type “identification at infinity” arguments or imposing Powell (1984, 1986a) type “sign” restrictions on the regression function to identify θ^* .

¹⁰If inference is not done conditional on the estimated value of K_0 (even though it is known), i.e., (4.3) is not taken into account and only (4.2) is used to estimate θ^* , then Theorem 4.1 shows that the resulting estimator of θ^* will be consistent but not asymptotically efficient. This apparent paradox where estimating a known parameter leads to a more efficient estimator of the parameter of interest arises quite frequently in the biased sampling or incomplete data literature; see, e.g., Cosslett (1981), Robins, Hsieh, and Newey (1995), and the references therein. As explained in Lancaster (1992), this happens when the conditionality principle is not followed in practice; i.e., the paradox disappears when inference is done conditional on the ancillary statistic.

So let $a(Z, K) = K + (1 - K)\mathbb{I}(Z \stackrel{elt}{<} c)$ and

$$\rho(Z, \theta, K) = \begin{bmatrix} \frac{g(Z, \theta)\mathbb{I}(Z \stackrel{elt}{\neq} c)}{a(Z, K)} \\ \frac{\mathbb{I}(Z \stackrel{elt}{\neq} c)}{a(Z, K)} - 1 \end{bmatrix} \stackrel{def}{=} \begin{bmatrix} \rho_1(Z, \theta, K) \\ \rho_2(Z, K) \end{bmatrix}_{(q+1) \times 1}, \quad (4.5)$$

where $\rho_1(Z, \theta, K) = g(Z, \theta)\mathbb{I}(Z \stackrel{elt}{\neq} c)/a(Z, K)$ and $\rho_2(Z, K) = \mathbb{I}(Z \stackrel{elt}{\neq} c)/a(Z, K) - 1$. Since K_0 is just identified by (4.4) and θ^* is globally identified in (2.1) by Assumption 4.1(iii),

$$\mathbb{E}_{f^*}\{g(Z^*, \theta^*)\} = 0 \iff \mathbb{E}_{f_e}\{\rho(Z, \theta^*, K_0)\} = 0. \quad (4.6)$$

Hence, efficient inference can be based on the latter moment condition without any loss of information.

Now let Z_1, \dots, Z_n denote iid random vectors with μ -density f_e . Using (4.6), θ^* and K_0 can be estimated by the usual two-step GMM as follows: First, use a preliminary consistent estimator $(\tilde{\theta}, \tilde{K})_{(p+1) \times 1} = \operatorname{argmin}_{(\theta, K) \in \Theta \times [0, 1]} \hat{\rho}'(\theta, K) \hat{\rho}(\theta, K)$, where $\hat{\rho}(\theta, K) = \sum_{j=1}^n \rho(Z_j, \theta, K)/n$, to construct the optimal weight matrix $\tilde{\Upsilon} = \sum_{j=1}^n \rho(Z_j, \tilde{\theta}, \tilde{K}) \rho'(Z_j, \tilde{\theta}, \tilde{K})/n$. Next, obtain the GMM estimator

$$(\hat{\theta}_{gmm}, \hat{K}_{gmm}) = \operatorname{argmin}_{(\theta, K) \in \Theta \times [0, 1]} \text{GMM}(\theta, K), \quad \text{where} \quad \text{GMM}(\theta, K) = \hat{\rho}'(\theta, K) \tilde{\Upsilon}^{-1} \hat{\rho}(\theta, K). \quad (4.7)$$

The following assumption, which is maintained throughout the paper, ensures that the GMM and EL estimators (described subsequently) are consistent and asymptotically normal. Let $\|\cdot\|$ denote the Euclidean norm and $B(\theta, \delta)$ an open ball with center θ and radius δ .

Assumption 4.1. (i) $K_0 \in (0, 1)$; (ii) Θ is compact; (iii) $\theta^* \in \operatorname{int}(\Theta)$ is the unique root of (2.1); (iv) $g(z, \theta)$ is continuous on Θ w.p.1; There exist $\eta > 0$ and $\delta > 0$ such that the following conditions hold: (v) $\mathbb{E}_{f^*}\{\sup_{\theta \in \Theta} \|g(z, \theta)\|^{2(1+\eta)}\} < \infty$; (vi) $g(z, \theta)$ is twice continuously differentiable on $B(\theta^*, \delta)$ w.p.1; (vii) $\mathbb{E}_{f^*}\{\sup_{\theta \in B(\theta^*, \delta)} \|\frac{\partial g(z, \theta)}{\partial \theta}\|\} < \infty$; (viii) $\mathbb{E}_{f^*}\{\sup_{\theta \in B(\theta^*, \delta)} \|\frac{\partial^2 g^{(i)}(z, \theta)}{\partial \theta^{(i)} \theta^{(k)}}\|\} < \infty$ for $i = 1, \dots, q$ and $j, k = 1, \dots, p$.

(i) rules out the uninteresting case (if $K_0 = 1$ the censoring or truncation problem vanishes). (iii) ensures that θ^* in (2.1) is globally identified. (i)–(viii) are used to prove the consistency and asymptotic normality of EL estimators as in Kitamura (1997) and Qin and Lawless (1994), although GMM estimators can be shown to be consistent and asymptotically normal under slightly weaker conditions; see, e.g., Newey and McFadden (1994).

It is well known from standard GMM theory that $\hat{\theta}_{gmm}$ and \hat{K}_{gmm} are jointly normal in large samples. In fact, letting $D_{q \times p} = \mathbb{E}_{f_e}\{\frac{\partial \rho_1(Z, \theta^*, K_0)}{\partial \theta}\}$, $V_{q \times q} = \mathbb{E}_{f_e}\{\rho_1(Z, \theta^*, K_0) \rho_1'(Z, \theta^*, K_0)\}$, $\gamma_{q \times 1} = \mathbb{E}_{f_e}\{\rho_1(Z, \theta^*, K_0) \rho_2(Z, K_0)\}$, $\delta = \mathbb{E}_{f_e}\{\rho_2^2(Z, \theta^*, K_0)\}$, and $\Omega = V - \gamma \gamma' / \delta$, we can show that

$$\mathbf{Theorem 4.1.} \quad \begin{bmatrix} n^{1/2}(\hat{\theta}_{gmm} - \theta^*) \\ n^{1/2}(\hat{K}_{gmm} - K_0) \end{bmatrix} \xrightarrow{d} N(0_{(p+1) \times 1}, \begin{bmatrix} (D' \Omega^{-1} D)^{-1} & 0_{p \times 1} \\ 0'_{p \times 1} & (1 - K_0)^2 / \delta \end{bmatrix}).$$

Since the limiting distribution is Gaussian with a diagonal variance-covariance matrix, $\hat{\theta}_{gmm}$ and \hat{K}_{gmm} are asymptotically independent. It follows that inference based on the marginal asymptotic distribution of $\hat{\theta}$ in Theorem 4.1 is equivalent to inference based on the asymptotic conditional

distribution of $\hat{\theta}|\hat{K}$. Furthermore, since it is well known that optimally weighted GMM estimators are efficient as the sample size goes to infinity (Chamberlain 1987), we also have that $\hat{\theta}_{gmm}$ is asymptotically efficient¹¹.

Before proceeding further, we now provide some intuition behind how transforming the moment condition allows us to impute the censored values. So, using (4.4), note that we can decompose

$$\begin{aligned} \mathbb{E}_{f_e}\{\rho_1(Z, \theta^*, K_0)\} &= \mathbb{E}_{f_e}\{g(Z, \theta^*)|Z <^{\text{elt}} c\}\Pr_{f_e}(Z <^{\text{elt}} c) \\ &\quad + \mathbb{E}_{f_e}\{g(Z, \theta^*)|(Z \neq^{\text{elt}} c) \cap (Z <^{\text{elt}} c)^{\sim}\}\Pr_{f_e}(\{Z <^{\text{elt}} c\}^{\sim}). \end{aligned} \quad (4.8)$$

Therefore, the moment condition in (4.2) can be expressed as a weighted sum of the best predictors of $g(Z^*, \theta^*)|(Z^*$ is uncensored) and $g(Z^*, \theta^*)|(Z^*$ is censored), with the weights being equal to the probability that Z^* is uncensored or censored, respectively. The estimators proposed in this paper use the enriched sample to automatically replace $g(Z^*, \theta^*)$ with its best predictor when observations are censored and then consistently and efficiently estimate these best predictors and selection probabilities; see Example 4.1 for a nice illustration.

Next, we show how to estimate θ^* by EL. So let p_j denote the probability mass placed at Z_j by a discrete distribution that has support on the realized observations Z_1, \dots, Z_n . For a fixed θ and K concentrate out the p_j 's by solving the nonparametric maximum likelihood problem $\max_{p_1, \dots, p_n} \sum_{j=1}^n \log p_j$ subject to the constraints that the p_j 's are nonnegative, $\sum_{j=1}^n p_j = 1$, and $\sum_{j=1}^n \rho(Z_j, \theta, K)p_j = 0$. If the convex hull of $\rho(Z_1, \theta, K), \dots, \rho(Z_n, \theta, K)$ contains the origin, then it is easy to verify that the solution to this problem is given by $\hat{p}_j(\theta, K) = n^{-1}\{1 + \lambda'(\theta, K)\rho(Z_j, \theta, K)\}^{-1}$ for $j = 1, \dots, n$, where the Lagrange multiplier $\lambda(\theta, K)$ satisfies $\sum_{j=1}^n \rho(Z_j, \theta, K)\{1 + \lambda'(\theta, K)\rho(Z_j, \theta, K)\}^{-1} = 0$. Letting $\text{EL}(\theta, K) = \sum_{j=1}^n \log \hat{p}_j(\theta, K) = -\sum_{j=1}^n \log\{1 + \lambda'(\theta, K)\rho(Z_j, \theta, K)\} - n \log n$, we can define the empirical likelihood estimators of θ^* and K_0 as $(\hat{\theta}_{el}, \hat{K}_{el}) = \text{argmax}_{(\theta, K) \in \Theta \times [0, 1]} \text{EL}(\theta, K)$, where

$$\text{EL}(\theta, K) = \sum_{j=1}^n \log \hat{p}_j(\theta, K) = -\sum_{j=1}^n \log\{1 + \lambda'(\theta, K)\rho(Z_j, \theta, K)\} - n \log n. \quad (4.9)$$

Consistency of $\hat{\theta}_{el}$ and \hat{K}_{el} follows from Kitamura (1997, Theorem 1). By standard EL theory as in Qin and Lawless (1994), EL and GMM estimators have the same asymptotic distribution. Hence, the EL estimator of θ^* is also asymptotically efficient. In finite samples the GMM and EL estimators will be different, though the two coincide if θ^* is just identified (i.e., $q = p$) because then the EL weights $\hat{p}_j(\theta, K) = 1/n$ for each j , θ , and K .

Although GMM and EL based statistical inference is asymptotically first order equivalent, recent research by Newey and Smith (2003) has shown that under certain regularity conditions EL has better second order properties than GMM. For instance, they show that, unlike GMM, the second

¹¹Readers interested in seeing this from first principles should examine the approach described in Tripathi (2003). In particular, the proof of Theorem 4.8 in Tripathi (2003) can be easily extended to show that the efficiency bound for estimating θ^* coincides with the asymptotic variance of $\hat{\theta}$ given in Theorem 4.1. For the sake of conciseness, we omit all efficiency bound calculations from this paper.

order bias of EL does not depend upon the number of moment conditions. This makes EL very attractive for estimating models with large q (e.g., panel data models with long time dimension) where GMM is known to perform poorly in small samples. As far as testing is concerned, Kitamura (2001) has demonstrated that an EL based specification test for (2.1) is asymptotically optimal in terms of a Hoeffding type large deviation criterion. Brown and Newey (2002) show that EL is intimately connected with the theory of efficient bootstrapping.

Another advantage of EL is that efficient estimators of the realized cdf $F_e(\xi) = \Pr_{f_e}(Z \leq \xi)$ and the target cdf $F^*(\xi)$ are easily obtained. So let $\hat{F}(\xi) = \sum_{j=1}^n \hat{p}_j(\hat{\theta}_{el}, \hat{K}_{el}) \mathbb{I}(Z_j \leq \xi)$ and $\hat{F}^*(\xi) = \sum_{j=1}^n \hat{p}_j(\hat{\theta}_{el}, \hat{K}_{el}) \mathbb{I}(Z_j \leq \xi) \mathbb{I}(Z_j \neq c) / a(Z_j, \hat{K}_{el})$, where the $\hat{p}_j(\hat{\theta}_{el}, \hat{K}_{el})$'s are the EL probability weights evaluated at $(\hat{\theta}_{el}, \hat{K}_{el})$. Then,

Theorem 4.2. $n^{1/2}\{\hat{F}(\xi) - F_e(\xi)\}$ is asymptotically normal with mean zero and variance

$$F_e(\xi)\{1 - F_e(\xi)\} - \mathbb{E}_{f_e}\{\tilde{g}'(Z)\mathbb{I}(Z \leq \xi)\}(\Omega^{-1} - \Omega^{-1}D(D'\Omega^{-1}D)^{-1}D'\Omega^{-1})\mathbb{E}_{f_e}\{\tilde{g}(Z)\mathbb{I}(Z \leq \xi)\},$$

where $\tilde{g}(Z) = \rho_1(Z, \theta^*, K_0) - (\gamma/\delta)\rho_2(Z, K_0)$.

Next, letting $b(\xi) = \mathbb{E}_{f_e}\{\mathbb{I}(Z \leq \xi)\mathbb{I}(Z \neq c)\mathbb{I}(Z < c)^\sim\} / \mathbb{E}_{f_e}\{\mathbb{I}(Z \neq c)\mathbb{I}(Z < c)^\sim\}$, define $m(Z, \xi) = \mathbb{I}(Z \leq \xi)\mathbb{I}(Z \neq c) - b(\xi)\rho_2(Z, K_0)a(Z, K_0)$ and $m_a(Z, \xi) = m(Z, \xi)/a(Z, K_0)$. Then,

Theorem 4.3. $n^{1/2}\{\hat{F}^*(\xi) - F^*(\xi)\}$ is asymptotically normal with mean zero and variance

$$\text{var}_{f_e}\{m_a(Z, \xi)\} - \mathbb{E}_{f_e}\{\tilde{g}'(Z)m_a(Z, \xi)\}(\Omega^{-1} - \Omega^{-1}D(D'\Omega^{-1}D)^{-1}D'\Omega^{-1})\mathbb{E}_{f_e}\{\tilde{g}(Z)m_a(Z, \xi)\}.$$

Theorems 4.2 and 4.3 show that imposing the overidentified model leads to an efficiency gain in estimating F_e and F^* . Asymptotic optimality of EL implies that $\hat{F}(\xi)$ and $\hat{F}^*(\xi)$ are also asymptotically efficient. Alternatively, this can be proved directly by using the approach of Tripathi (2003) to first calculate the efficiency bounds for estimating $F_e(\xi)$ and $F^*(\xi)$ and then comparing these bounds with the asymptotic variances in Theorems 4.2 and 4.3.

For the remainder of Section 4, let $(\hat{\theta}, \hat{K})$ denote the GMM or EL estimator of (θ^*, K_0) . The asymptotic variances of $\hat{\theta}$, $\hat{F}(\xi)$, and $\hat{F}^*(\xi)$ can be estimated in the obvious manner by replacing D and Ω with consistent estimators $\hat{D} = \sum_{j=1}^n \frac{\partial \rho_1(Z_j, \hat{\theta}, \hat{K})}{\partial \theta} / n$ and $\hat{\Omega} = \hat{V} - \hat{\gamma}\hat{\gamma}' / \hat{\delta}$, respectively, where $\hat{V} = \sum_{j=1}^n \rho_1(Z_j, \hat{\theta}, \hat{K})\rho_1'(Z_j, \hat{\theta}, \hat{K}) / n$, $\hat{\gamma} = \sum_{j=1}^n \rho_1(Z_j, \hat{\theta}, \hat{K})\rho_2(Z_j, \hat{K}) / n$, and $\hat{\delta} = \sum_{j=1}^n \rho_2^2(Z_j, \hat{K}) / n$.

Example 4.1 (Example 3.1 contd.). Here $\rho_1(Z, \theta, K) = (Z - \theta)\mathbb{I}(Z \neq c) / a(Z, K)$ and, since there are no overidentifying restrictions, $\hat{\theta}$ and \hat{K} solve $\sum_{j=1}^n \rho(Z_j, \hat{\theta}, \hat{K}) = 0$; i.e.,

$$\hat{\theta} = n^{-1} \sum_{j=1}^n \frac{Z_j \mathbb{I}(Z_j \neq c)}{a(Z_j, \hat{K})} \quad \text{and} \quad \hat{K} = \frac{\sum_{j=1}^n \mathbb{I}(Z_j \neq c)\mathbb{I}(Z_j < c)^\sim}{\sum_{j=1}^n \mathbb{I}(Z_j < c)^\sim}. \quad (4.10)$$

To gain further insight into $\hat{\theta}$, notice that for $d = 1$ we can write

$$\hat{\theta} = n^{-1} \sum_{j=1}^n \mathbb{I}(Z_j < c) \times \frac{\sum_{j=1}^n Z_j \mathbb{I}(Z_j < c)}{\sum_{j=1}^n \mathbb{I}(Z_j < c)} + n^{-1} \sum_{j=1}^n \mathbb{I}(Z_j \geq c) \times \frac{\sum_{j=1}^n Z_j \mathbb{I}(Z_j > c)}{\sum_{j=1}^n \mathbb{I}(Z_j > c)}.$$

In light of (4.8), it comes as no surprise that $\hat{\theta}$ is a linear combination of the sample means of uncensored and censored observations in the enriched dataset with the weights being the fraction of uncensored and censored observations in the enriched sample.

By Theorem 4.1, $n^{1/2}(\hat{\theta} - \theta^*)$ converges in distribution to a normal random variable with mean zero and variance Ω . Since there are no overidentifying restrictions in this example (and the next one), $\hat{F}(\xi) = n^{-1} \sum_{j=1}^n \mathbb{I}(Z_j \stackrel{elt}{\leq} \xi)$ is just the empirical cdf of the realized observations and $\hat{F}^*(\xi) = n^{-1} \sum_{j=1}^n \mathbb{I}(Z_j \stackrel{elt}{\leq} \xi) \mathbb{I}(Z_j \neq c) / a(Z_j, \hat{K})$. By Theorem 4.2, or by a direct application of the central limit theorem, the asymptotic variance of $n^{1/2}\{\hat{F}(\xi) - F_e(\xi)\}$ is simply $F_e(\xi)\{1 - F_e(\xi)\}$. Similarly, by Theorem 4.3, $n^{1/2}\{\hat{F}^*(\xi) - F^*(\xi)\}$ has asymptotic variance $var_{f_e}\{m_a(Z, \xi)\}$. \square

Example 4.2 (Example 3.2 contd.). Here, $\rho_1(Z, \theta, K) = X(Y - X'\theta) \mathbb{I}(Z \neq c) / a(Z, K)$. Hence, $\hat{\theta} = (\sum_{j=1}^n \hat{X}_j X_j')^{-1} (\sum_{j=1}^n \hat{X}_j Y_j)$, where $\hat{X} = X \mathbb{I}(Z \neq c) / a(Z, \hat{K})$ and \hat{K} is given in (4.10). Notice that this tantamounts to replacing the original regressors in Example 3.2 with new instruments \hat{X} . If censoring is purely endogenous or purely exogenous, then $a(Z, K) = K + (1 - K) \mathbb{I}(Y_j < c^{(1)})$ or $a(Z, K) = K + (1 - K) \mathbb{I}(X_j \stackrel{elt}{<} c^{-(1)})$, respectively, and the expression for $\hat{\theta}$ simplifies. \square

Example 4.3 (Censored linear regression with endogenous regressors). Let $Y^* = X^{*'}\theta^* + \varepsilon$ such that some or all of the regressors are correlated with ε . We have a $q \times 1$ vector of instrumental variables W^* that are uncorrelated with ε ; i.e., W^* satisfies the moment condition $\mathbb{E}_{f^*}\{W^* \varepsilon\} = 0$. Let $W^* = (X_1^*, \tilde{W}^*)$, where X_1^* denotes the $p_1 \times 1$ vector of exogenous coordinates of X^* and \tilde{W}^* the $(q - p_1) \times 1$ vector of instruments for the endogenous coordinates of X^* . Hence, in this example, $Z^* = (Y^*, X^*, \tilde{W}^*)$ and $g(Z^*, \theta^*) = W^*(Y^* - X^{*'}\theta^*)$. If the dependent variable, regressors, and instruments are all censored, then $\rho_1(Z, \theta, K) = W(Y - X'\theta) \mathbb{I}(Z \neq c) / a(Z, K)$. On the other hand, the endogenous tobit model where only Y^* is censored and X^* is endogenous is very important for applications and follows by letting $c^{-(1)} = (\infty, \dots, \infty)$ so that $\rho_1(Z, \theta, K) = W(Y - X'\theta) \mathbb{I}(Y \neq c^{(1)}) / a(Y, K)$, where $a(Y, K) = K + (1 - K) \mathbb{I}(Y < c^{(1)})$. In either case, θ^* can be estimated by GMM or EL as described earlier and the asymptotic distribution of $\hat{\theta}$ follows readily from Theorem 4.1. \square

Example 4.4 (Censoring and IV). Suppose $Y_1^* = X_1^{*'}\theta_1^* + \varepsilon_1$ and $Y_2^* = X_2^{*'}\theta_2^* + \varepsilon_2$, where $\varepsilon = (\varepsilon_1, \varepsilon_2)_{2 \times 1}$ satisfies the conditional moment restriction $\mathbb{E}_{(Y_1^*, Y_2^*)|X^*}(\varepsilon|X^*) = 0$ w.p.1 and X^* denotes a vector containing the exogenous coordinates of $(X_1^*, X_2^*)_{(p_1+p_2) \times 1}$ and other instruments. Hence, $\mathbb{E}_{f^*}\{A(X^*) \begin{bmatrix} Y_1^* - X_1^{*'}\theta_1^* \\ Y_2^* - X_2^{*'}\theta_2^* \end{bmatrix}\} = 0$, where $A(X^*)$ is a $q \times 2$ matrix of instrumental variables such that $q \geq p_1 + p_2$. If $Z^* = (Y_1^*, Y_2^*, X^*)$ is censored, then (4.6) can be used to estimate $\theta^* = (\theta_1^*, \theta_2^*)$ by GMM or EL as proposed earlier. Although this simultaneous equations model has been studied before, see, e.g., Smith and Blundell (1986) and Blundell and Smith (1993), the treatment here is

more general because we do not assume that ε is Gaussian and allow for the possibility that besides Y_1^* and Y_2^* the instruments may also be censored. Censoring of $Y^* = (Y_1^*, Y_2^*)$ alone means that $c^{-(1,2)} = (\infty, \dots, \infty)$. Hence, $\rho_1(Z, \theta, K) = A(X) \begin{bmatrix} Y_1 - X_1' \theta_1 \\ Y_2 - X_2' \theta_2 \end{bmatrix} \mathbb{I}(Y_1^* \neq c^{(1)}, Y_2^* \neq c^{(2)}) / a(Y, K)$, where $a(Y, K) = K + (1 - K) \mathbb{I}(Y_1 < c^{(1)}, Y_2 < c^{(2)})$, and θ^* can be estimated by GMM or EL as before. \square

Example 4.5. Sometimes we may possess auxiliary information about a feature of the target density; e.g., we may know beforehand that the mean of the target population is zero. In general, suppose it is known a priori that $\mathbb{E}_{f^*}\{m(Z^*)\} = 0$, where m is a vector of known functions. Moment based auxiliary information about f^* can be easily incorporated in our framework by first stacking $g(Z^*, \theta^*)$ and $m(Z^*)$ to produce an augmented vector of moment conditions and then proceeding as before. These types of models, which are a special case of the general unconditional moment restrictions model examined in this paper, have been investigated by Imbens and Lancaster (1994), Hellerstein and Imbens (1999), and Nevo (2003). However, Imbens and Lancaster (1994) and Hellerstein and Imbens (1999) assume that Z^* is fully observed. Nevo (2003) allows Z^* to be entirely missing (due to attrition) but not censored. He also restricts attention to the case where the parameter of interest is just identified. In addition, he assumes that the selection probability is known up to a finite dimensional parameter and imposes an identification condition that rules out truncated Z^* 's as well. In contrast, we allow (2.1) to be overidentified and the selection probabilities for censoring or truncation of Z^* to be fully unknown. Although Nevo also uses a supplementary sample to estimate finite dimensional parameters, unlike us he does not consider identification and estimation of the target or realized cdf's. None of these papers discuss efficient estimation of the target and realized cdf's. \square

4.2. Hypothesis tests and confidence regions. Suppose we want to test the parametric restriction $\tilde{H}_0 : R(\theta^*) = 0$ against the alternative that it is false, where R is a $\bar{r} \times 1$ vector of twice continuously differentiable functions such that $\frac{\partial R(\theta^*)}{\partial \theta}$ has rank $\bar{r} \leq p$. Since $n^{1/2}(\hat{\theta} - \theta^*) \xrightarrow{d} N(0, (D' \Omega^{-1} D)^{-1})$, the Wald statistic $W = nR'(\hat{\theta}) \left\{ \frac{\partial R(\hat{\theta})}{\partial \theta} (\hat{D}' \hat{\Omega}^{-1} \hat{D})^{-1} \frac{\partial R'(\hat{\theta})}{\partial \theta} \right\}^{-1} R(\hat{\theta}) \xrightarrow{d} \chi_{\bar{r}}^2$. Hence, a test for \tilde{H}_0 can be based upon W .

Another alternative is to use a distance metric test. So let $\bar{\theta}$ and \bar{K} denote the GMM and EL estimators under \tilde{H}_0 ; i.e., $(\bar{\theta}_{gmm}, \bar{K}_{gmm}) = \operatorname{argmin}_{\{(\theta, K) \in \Theta \times [0, 1] : R(\theta) = 0\}} \operatorname{GMM}(\theta, K)$ and $(\bar{\theta}_{el}, \bar{K}_{el}) = \operatorname{argmax}_{\{(\theta, K) \in \Theta \times [0, 1] : R(\theta) = 0\}} \operatorname{EL}(\theta, K)$. Next, let $DM = n \{ \operatorname{GMM}(\bar{\theta}_{gmm}, \bar{K}_{gmm}) - \operatorname{GMM}(\hat{\theta}_{gmm}, \hat{K}_{gmm}) \}$ and $LR = 2 \{ \operatorname{EL}(\hat{\theta}_{el}, \hat{K}_{el}) - \operatorname{EL}(\bar{\theta}_{el}, \bar{K}_{el}) \}$, where DM denotes the GMM based distance metric statistic and LR the EL based likelihood ratio test statistic. A test for \tilde{H}_0 can also be based upon DM or LR . To obtain the critical values for these tests we can use Newey and McFadden (1994, Theorem 9.2) and Qin and Lawless (1994, Theorem 2) to see that $DM \xrightarrow{d} \chi_{\bar{r}}^2$ and $LR \xrightarrow{d} \chi_{\bar{r}}^2$ under \tilde{H}_0 .

Since W , DM , and LR are asymptotically equivalent, the decision to use a particular test statistic in practice usually depends upon computational and other considerations; e.g., although all three statistics can be inverted to obtain asymptotically valid confidence regions, the DM and LR based regions are invariant to the formulation of \tilde{H}_0 and automatically satisfy natural range restrictions. Furthermore, unlike W and DM , the likelihood ratio statistic LR is internally studentized; i.e., it does

not require preliminary estimation of any variance terms. This guarantees that confidence regions based on LR are also invariant to nonsingular transformations of the moment conditions.

4.3. Specification tests. For the remainder of this section, assume that $q > p$. Since inference based on the estimated θ^* is sensible only if (2.1) is true, it is important to test H_0 against the alternative that it is false. In this section, we describe two ways of testing H_0 . The first approach is easy: GMM theory tells us that $n\text{GMM}(\hat{\theta}_{gmm}, \hat{K}_{gmm}) \xrightarrow{d} \chi_{q-p}^2$ under H_0 . Therefore, a test for overidentifying restrictions (usually called the J -test) in (2.1) can be based on this result.

An EL based specification test for H_0 can also be developed. Besides being internally studentized and invariant to nonsingular and algebraic transformations of the moment conditions, this test has been shown by Kitamura (2001) to be optimal in terms of a large deviations criterion.

So let $\hat{\theta}$ and \hat{K} denote $n^{1/2}$ -consistent preliminary estimators of θ^* and K_0 ; e.g., $\hat{\theta}$ and \hat{K} can be the GMM or EL estimators defined previously. The restricted (i.e., under H_0) EL can be written as $\text{EL}^r = \sum_{j=1}^n \log \hat{\rho}_j(\hat{\theta}, \hat{K})$. Next, consider the unrestricted problem where the model is not imposed. It is well known that the nonparametric maximum likelihood estimator of f_e in the absence of any auxiliary information puts mass $1/n$ at each realized observation and is zero elsewhere. Therefore, the unrestricted nonparametric likelihood is given by $\text{EL}^{ur} = -n \log n$. Now let $\text{ELR} = 2(\text{EL}^{ur} - \text{EL}^r) = 2 \sum_{j=1}^n \log\{1 + \lambda'(\hat{\theta}, \hat{K})\rho(Z_j, \hat{\theta}, \hat{K})\}$. Then ELR can be regarded as an analog of the usual parametric likelihood ratio test statistic; i.e., H_0 is rejected if ELR is large enough. By Qin and Lawless (1994, Corollary 4), $\text{ELR} \xrightarrow{d} \chi_{q-p}^2$ under H_0 . Hence, critical values for ELR are easily obtained.

5. INFERENCE WITH TRUNCATED DATA

We now show how to do efficient inference using truncated data. As before, we begin by demonstrating that f^* is identified if the realized observations are drawn from the enriched mixture density defined in (2.5).

5.1. Identification and efficient estimation. Let $b^* = \int_T f^*(z) d\mu^*$ denote the probability that Z^* is observed. Throughout the paper we assume that $b^* \in (0, 1)$ is unknown. Since K_0 is known and b^* can be explicitly written in terms of the realized enriched density alone as

$$b^* = \frac{\mathbb{E}_{f_e}[\mathbb{I}(Z \in T)] - (1 - K_0)}{K_0}, \quad (5.1)$$

it follows that b^* is identified. Next, by (2.4) and (2.5),

$$f^*(z) = \frac{f_e(z)}{K_0 + (1 - K_0)\mathbb{I}(z \in T)/b^*} \quad (5.2)$$

for all $z \in \mathbb{R}^d$, where b^* is given by (5.1). Hence, f^* is also identified.

Estimation is also straightforward, although now there is an additional parameter b^* to estimate. So let $a(Z, b) = K_0 + (1 - K_0)\mathbb{I}(Z \in T)/b$ and define¹²

$$\rho(Z, \theta, b) = \begin{bmatrix} \frac{g(Z, \theta)}{a(Z, b)} \\ \frac{\mathbb{I}(Z \in T) - (1 - K_0)}{K_0} - b \end{bmatrix} \stackrel{def}{=} \begin{bmatrix} \rho_1(Z, \theta, b) \\ \rho_2(Z, b) \end{bmatrix}_{(q+1) \times 1}, \quad (5.3)$$

where $\rho_1(Z, \theta, b) = g(Z, \theta)/a(Z, b)$ and $\rho_2(Z, b) = \{\mathbb{I}(Z \in T) - (1 - K_0)\}/K_0 - b$. Since, by (5.2),

$$\mathbb{E}_{f^*}\{g(Z^*, \theta^*)\} = 0 \iff \mathbb{E}_{f_e}\{\rho(Z, \theta^*, b^*)\} = 0, \quad (5.4)$$

θ^* and b^* can be jointly and efficiently estimated by using the latter moment condition. In particular, the GMM and EL estimators of θ^* and b^* are given by $(\hat{\theta}_{gmm}, \hat{b}_{gmm}) = \operatorname{argmin}_{(\theta, b) \in \Theta \times [0, 1]} \text{GMM}(\theta, b)$ and $(\hat{\theta}_{el}, \hat{b}_{el}) = \operatorname{argmax}_{(\theta, b) \in \Theta \times [0, 1]} \text{EL}(\theta, b)$, respectively, where the objective functions are obtained from (4.7) and (4.9) by replacing K with b .

The logic behind the imputation mechanism becomes clear upon observing that

$$\mathbb{E}_{f_e}\{\rho_1(Z, \theta^*, b^*)\} = b^* \mathbb{E}_{f_e}\{g(Z, \theta^*) | Z \in T\} + (1 - b^*) \mathbb{E}_{f_e}\{g(Z, \theta^*) | Z \notin T\}; \quad (5.5)$$

i.e., the transformed moment condition in (5.4) is just a combination of the best predictors of $g(Z^*, \theta^*)$ (Z^* is not truncated) and $g(Z^*, \theta^*)$ (Z^* is truncated) weighted by probabilities of the corresponding events. As in the case of censoring, the GMM or EL estimators we propose automatically carry out this imputation in the enriched sample to efficiently estimate $\hat{\theta}$ and \hat{b} .

Let $g_a(Z, \theta^*) = g(Z, \theta^*)/a(Z, b^*)$, $D = \mathbb{E}_{f_e}\{\frac{\partial g_a(Z, \theta^*)}{\partial \theta}\}$, and $\Omega = \operatorname{var}_{f_e}\{\tilde{g}_a(Z, \theta^*)\}$, where $\tilde{g}_a(Z, \theta^*) = \tilde{g}(Z, \theta^*)/a(Z, b^*)$ and $\tilde{g}(Z, \theta^*) = g(Z, \theta^*) + K_0^{-1}(1 - K_0)b^{*-2}\mathbb{I}(Z \in T)\mathbb{E}_{f_e}\{g_a(Z, \theta^*)\mathbb{I}(Z \in T)\}$. Also, let $(\hat{\theta}, \hat{b})$ denote the GMM or EL estimator of (θ^*, b^*) for the remainder of the paper. Then, letting $\Sigma_{11} = (D'\Omega^{-1}D)^{-1}$, $G = \mathbb{E}_{f_e}\{g_a(Z, \theta^*)\mathbb{I}(Z \in T)\}$, $\Sigma_{12} = -(K_0b^*)^{-1}(K_0b^* + 1 - K_0)(D'\Omega^{-1}D)^{-1}D'\Omega^{-1}G$, and $\Sigma_{22} = K_0^{-2}\operatorname{var}_{f_e}\{\mathbb{I}(Z \in T)\} - (K_0b^*)^{-2}(K_0b^* + 1 - K_0)^2G'(\Omega^{-1} - \Omega^{-1}D(D'\Omega^{-1}D)^{-1}D'\Omega^{-1})G$, we have that

Theorem 5.1. $\begin{bmatrix} n^{1/2}(\hat{\theta} - \theta^*) \\ n^{1/2}(\hat{b} - b^*) \end{bmatrix} \xrightarrow{d} \text{N}(0, \begin{bmatrix} \Sigma_{11} & \Sigma_{12} \\ \Sigma_{12}' & \Sigma_{22} \end{bmatrix})$.

It is intuitively clear from (5.4) that $\hat{\theta}$ and \hat{b} are asymptotically efficient¹³. Their asymptotic variances can be estimated by replacing Σ_{ij} 's with their consistent estimators constructed, say, using $\hat{D} = n^{-1} \sum_{j=1}^n \frac{\partial g_a(Z_j, \hat{\theta})}{\partial \theta}$ and $\hat{\Omega} = n^{-1} \sum_{j=1}^n \hat{g}_a(Z_j, \hat{\theta}) \hat{g}_a'(Z_j, \hat{\theta}) - n^{-1} \sum_{j=1}^n \hat{g}_a(Z_j, \hat{\theta}) n^{-1} \sum_{j=1}^n \hat{g}_a'(Z_j, \hat{\theta})$, where $\hat{a} \stackrel{def}{=} a(Z, \hat{b})$, $\hat{g}(Z, \theta) = g(Z, \theta) + K_0^{-1}(1 - K_0)\hat{b}^{-2}\mathbb{I}(Z \in T)\hat{G}$ and $\hat{G} = n^{-1} \sum_{j=1}^n g_a(Z_j, \hat{\theta})\mathbb{I}(Z_j \in T)$. Of course, $\operatorname{var}_{f_e}\{\mathbb{I}(Z \in T)\}$ is estimated by $(n_T/n)(1 - n_T/n)$, where $n_T = \sum_{j=1}^n \mathbb{I}\{Z_j \in T\}$ is the number of observations falling in T .

Next, we efficiently estimate $F_e(\xi)$ and $F^*(\xi)$. So let $\hat{F}(\xi) = \sum_{j=1}^n \hat{p}_j(\hat{\theta}_{el}, \hat{b}_{el})\mathbb{I}(Z_j \leq \xi)$ and $\hat{F}^*(\xi) = \sum_{j=1}^n \hat{p}_j(\hat{\theta}_{el}, \hat{b}_{el})\mathbb{I}(Z_j \leq \xi)/a(Z_j, \hat{b}_{el})$, where $\hat{p}_j(\hat{\theta}_{el}, \hat{b}_{el})$ are the EL probability weights evaluated at the EL estimators $(\hat{\theta}_{el}, \hat{b}_{el})$. Then,

¹²By (5.1), the second moment condition in $\rho(Z, \theta, b)$ is just the identification condition for b^* .

¹³More rigorously, as in the case of censoring, the proof of Theorem 4.8 in Tripathi (2003) can be easily extended to show that the efficiency bounds for estimating (θ^*, b^*) coincide with the asymptotic variances of $(\hat{\theta}, \hat{b})$ in Theorem 5.1.

Theorem 5.2. $n^{1/2}\{\hat{F}(\xi) - F_e(\xi)\}$ is asymptotically normal with mean zero and variance

$$F_e(\xi)\{1 - F_e(\xi)\} - \text{cov}_{f_e}\{\tilde{g}'_a(Z, \theta^*), \mathbb{I}(Z \leq \xi)\}(\Omega^{-1} - \Omega^{-1}D(D'\Omega^{-1}D)^{-1}D'\Omega^{-1})\text{cov}_{f_e}\{\tilde{g}_a(Z, \theta^*), \mathbb{I}(Z \leq \xi)\}^{elt}.$$

Defining $m(Z, \xi) = \mathbb{I}(Z \leq \xi) + [(1 - K_0)/(K_0 b^{*2})]\mathbb{I}(Z \in T)\mathbb{E}_{f_e}\{\mathbb{I}(Z \leq \xi)\mathbb{I}(Z \in T)/a(Z, b^*)\}$ and $m_a(Z, \xi) = m(Z, \xi)/a(Z, b^*)$, we can also show that

Theorem 5.3. $n^{1/2}\{\hat{F}^*(\xi) - F^*(\xi)\}$ is asymptotically normal with mean zero and variance

$$\text{var}_{f_e}\{m_a(Z, \xi)\} - \text{cov}_{f_e}\{\tilde{g}'_a(Z, \theta^*), m_a(Z, \xi)\}(\Omega^{-1} - \Omega^{-1}D(D'\Omega^{-1}D)^{-1}D'\Omega^{-1})\text{cov}_{f_e}\{\tilde{g}_a(Z, \theta^*), m_a(Z, \xi)\}.$$

As before, Theorems 5.2 and 5.3 show that if the model is overidentified, then imposing it leads to an efficiency gain in estimating F_e and F^* . Asymptotic optimality of EL implies that $\hat{F}(\xi)$ and $\hat{F}^*(\xi)$ are also asymptotically efficient.

Example 5.1 (Example 3.3 contd.). Since here $\rho_1(Z, \theta, b) = (Z - \theta)/a(Z, b)$ and there are no overidentifying restrictions, $\hat{\theta}$ and \hat{b} are obtained by solving $\sum_{j=1}^n \rho(Z_j, \hat{\theta}, \hat{b}) = 0$. Therefore, $\hat{b} = [n_T/n - (1 - K_0)]/K_0$ and $\hat{\theta} = n^{-1} \sum_{j=1}^n Z_j/a(Z_j, \hat{b})$ since $\sum_{j=1}^n 1/a(Z_j, \hat{b}) = n$. A little algebra shows that

$$\hat{\theta} = \hat{b} \times \frac{\sum_{j=1}^n Z_j \mathbb{I}(Z_j \in T)}{\sum_{j=1}^n \mathbb{I}(Z_j \in T)} + (1 - \hat{b}) \times \frac{\sum_{j=1}^n Z_j \mathbb{I}(Z_j \notin T)}{\sum_{j=1}^n \mathbb{I}(Z_j \notin T)},$$

which is what we would expect from (5.5). By Theorem 5.1, $n^{1/2}(\hat{b} - b^*)$ has asymptotic variance $\text{var}_{f_e}\{\mathbb{I}(Z \in T)\}/K_0^2$. The asymptotic variance of $n^{1/2}(\hat{\theta} - \theta^*)$ can be similarly obtained. The absence of overidentifying restrictions also implies that $\hat{F}(\xi) = n^{-1} \sum_{j=1}^n \mathbb{I}(Z_j \leq \xi)$ and $\hat{F}^*(\xi) = n^{-1} \sum_{j=1}^n \mathbb{I}(Z_j \leq \xi)/a(Z_j, \hat{b}_{el})$ in this example and the next one. Therefore, by Theorem 5.2 or directly by the central limit theorem, $n^{1/2}\{\hat{F}(\xi) - F_e(\xi)\}$ is asymptotically normal with mean zero and variance $F_e(\xi)\{1 - F_e(\xi)\}$. Similarly, by Theorem 5.3, $n^{1/2}\{\hat{F}^*(\xi) - F^*(\xi)\}$ is asymptotically normal with mean zero and variance $\text{var}_{f_e}\{m_a(Z, \xi)\}$. \square

Example 5.2 (Example 3.4 contd.). Since $\rho_1(Z, \theta, b) = X(Y - X'\theta)/a(Y, X, b)$ and $\rho_2(Z, b) = \{\mathbb{I}(Y \in T_1, X \in T_2) - (1 - K_0)/K_0\} - b$, where $a(Y, X, b) = K_0 + (1 - K_0)\mathbb{I}(Y \in T_1, X \in T_2)/b$, we have $\hat{\theta} = \{\sum_{j=1}^n X_j X'_j/a(Y_j, X_j, \hat{b})\}^{-1} \{\sum_{j=1}^n X_j Y_j/a(Y_j, X_j, \hat{b})\}$ and \hat{b} as given in the previous example.

Hence, $n^{1/2}(\hat{\theta} - \theta^*) \xrightarrow{d} N(0, D^{-1}\Omega D^{-1})$ by Theorem 5.1, where $D = -\mathbb{E}_{f_e}\{XX'/a(Y, X, b^*)\}$ and

$$\Omega = \text{var}_{f_e}\left\{\frac{X(Y - X'\theta^*)}{a(Y, X, b^*)} + \frac{(1 - K_0)\mathbb{I}(Y \in T_1, X \in T_2)}{K_0 b^{*2}} \frac{X(Y - X'\theta^*)\mathbb{I}(Y \in T_1, X \in T_2)}{a(Y, X, b^*)}\right\}.$$

If truncation is purely endogenous, then

$$\hat{\theta} = \left\{\sum_{j=1}^n \frac{X_j X'_j}{K_0 + (1 - K_0)\mathbb{I}(Y_j \in T_1)/\hat{b}}\right\}^{-1} \left\{\sum_{j=1}^n \frac{X_j Y_j}{K_0 + (1 - K_0)\mathbb{I}(Y_j \in T_1)/\hat{b}}\right\}.$$

Similarly, under pure exogenous truncation,

$$\hat{\theta} = \left\{\sum_{j=1}^n \frac{X_j X'_j}{K_0 + (1 - K_0)\mathbb{I}(X_j \in T_2)/\hat{b}}\right\}^{-1} \left\{\sum_{j=1}^n \frac{X_j Y_j}{K_0 + (1 - K_0)\mathbb{I}(X_j \in T_2)/\hat{b}}\right\}.$$

The asymptotic variances of $\hat{\theta}$ under pure endogenous or exogenous truncation can be obtained by setting $T_2 = \mathbb{R}^p$ or $T_1 = \mathbb{R}$, respectively, in the expressions for D and Ω given above. \square

Example 5.3 (Truncated linear regression with endogenous regressors). Consider the setup of Example 4.3, but now suppose that instead of being censored the target variable Z^* is truncated outside $T_1 \times T_2 \times T_3$. If the dependent variable, regressors, and instruments are all subject to truncation, then $\rho_1(Z, \theta, b) = W(Y - X'\theta)/a(Y, X, \tilde{W}, b)$ and $\rho_2(Z, b) = \{\mathbb{I}(Y \in T_1, X \in T_2, \tilde{W} \in T_3) - (1 - K_0)\}/K_0 - b$, where $a(Y, X, \tilde{W}, b) = K_0 + (1 - K_0)\mathbb{I}(Y \in T_1, X \in T_2, \tilde{W} \in T_3)/b$. Hence, θ^* and b^* can be estimated by GMM or EL as described earlier. Under pure endogenous truncation (i.e., $T_2 = \mathbb{R}^p$ and $T_3 = \mathbb{R}^{q-p_1}$), it follows from Theorem 5.1 that $n^{1/2}(\hat{\theta} - \theta^*)$ is asymptotically normal with mean zero and variance $(D'\Omega^{-1}D)^{-1}$, where $D = -\mathbb{E}_{f_e}\{WX'/a(Y, b^*)\}$,

$$\Omega = \text{var}_{f_e} \left\{ \frac{W(Y - X'\theta^*)}{a(Y, b^*)} + \frac{(1 - K_0)\mathbb{I}(Y \in T_1)}{K_0 b^{*2}} \frac{\mathbb{I}(Y \in T_1)}{a(Y, b^*)} \mathbb{E}_{f_e} \left[\frac{W(Y - X'\theta^*)\mathbb{I}(Y \in T_1)}{a(Y, b^*)} \right] \right\},$$

and $a(Y, b^*) = K_0 + (1 - K_0)\mathbb{I}(Y \in T_1)/b^*$. A similar result holds for pure exogenous truncation. \square

Example 5.4 (Truncation and IV). Again, consider the simultaneous equations model of Example 4.4, but now assume that Z^* is truncated instead of being censored. To further simplify the exposition, suppose that only Y_1^* and Y_2^* are truncated outside T_1 and T_2 , respectively. θ^* and b^* can then be estimated by GMM or EL upon noting that $\rho_1(Z, \theta, b) = A(X) \begin{bmatrix} Y_1 - X_1'\theta_1 \\ Y_2 - X_2'\theta_2 \end{bmatrix} / a(Y, b)$ and $\rho_2(Z, b) = \{\mathbb{I}(Y_1 \in T_1, Y_2 \in T_2) - (1 - K_0)\}/K_0 - b$, where $a(Y, b) = K_0 + (1 - K_0)\mathbb{I}(Y_1 \in T_1, Y_2 \in T_2)/b$. \square

5.2. Hypothesis and specification tests. Hypotheses of the form $R(\theta^*) = 0$ can be tested using the Wald, DM, or LR statistics as described in Section 4.2 by basing the test on $\rho(Z, \theta^*, b^*)$. In each case the test statistic is asymptotically distributed as a χ^2_r random variable under the null hypothesis. Similarly, if $q > p$ then a test for overidentifying moment restrictions can be done using the GMM based J -statistic or the ELR statistic on the moment vector $\rho(Z, \theta^*, b^*)$. The details are analogous to those in Section 4.3. In either case, the test statistic has an asymptotic χ^2_{q-p} distribution under H_0 .

6. APPLICATION

Our application studies the effects of changes in compulsory schooling laws on age at first marriage. While the primary purpose of the application is to demonstrate the methodology developed in this paper, this is also a topic of some substantive importance. Our data are 1% samples from the Public Use Files of the U.S. Census of Population for the years 1960, 1970, and 1980.

Understanding the determinants of age at first marriage is considered to be important for several reasons. In recent years, age at first marriage has risen. Much literature suggests that a rising age at first marriage may be socially undesirable because marriage may encourage good behavior and outcomes. For example, Akerlof (1998) provides evidence that marriage has a beneficial effect on male behavior, leading to a decrease in socially undesirable activities such as alcoholism, drug abuse, and violence. Also, Korenman and Neumark (1991) find that in the cross-section, married men earn about 11% more than observationally equivalent unmarried men. When they utilize panel data and

estimate a fixed effects model, the marriage effect is about $2/3$ the size of the cross-sectional estimate. Thus, it appears that there is a direct effect of being married on male earnings. However, in other work, they find that marriage reduces female participation and does not positively impact their wage rates (Korenman and Neumark 1992). Second, there is a great deal of concern about the effects of out-of-wedlock childbearing on single parents and their children. If rising age at first marriage is not accompanied by postponed childbearing, this problem becomes more severe. Relatedly, it has long been known, see, e.g., Coale (1971), that age at first marriage is an important determinant of fertility. However, rising age at first marriage may also have socially beneficial effects (Goldin and Katz 2002) because they have been linked to greater opportunities for young people, especially women, to obtain education and develop a professional career.

Theoretically, the effects of increased education on age of marriage are unclear. Koball (1998) describes the “economic provider” hypothesis that men are less likely to marry until they are securely employed. Because more education leads to higher earnings, it may lead to earlier marriage through this channel. The “adult transition” hypothesis proposes that events that delay the transition to adulthood will also delay marriage. More education will tend to delay marriage through this channel. Empirically, there is a positive relationship between education and age of marriage and rising education may be related to increased age at first marriage in recent decades. However, the correlation between education and age at first marriage may reflect the fact that young people with low ability and poor labor market prospects choose both to marry early and to drop out of school early rather than a causal relationship between education and age at first marriage. One way to examine this issue is to look at the effects of changes in policy that led to increased education. In particular, we study whether increased mandatory educational attainment (through compulsory schooling legislation) encourages people to defer marriage. If so, these factors should be considered when evaluating the benefits of this type of legislation.

We use variation in compulsory schooling laws across states and over time. Changes in these laws had a significant impact on education and indeed have been used as instruments for education in other contexts by Acemoglu and Angrist (2001), Lochner and Moretti (2004), and Lleras-Muney (2002). Since the history of compulsory schooling laws in the U.S. is by now well documented (see, in particular, Lleras-Muney (2001) and Goldin and Katz (2003)), we will not describe them in great detail here. Essentially, there were five possible restrictions on educational attendance: (i) maximum age by which a child must be enrolled, (ii) minimum age at which a child may drop out, (iii) minimum years of schooling before dropping out, (iv) minimum age for a work permit, and (v) minimum schooling required for a work permit. In the years relevant to our sample, 1939 to 1958, states changed compulsory attendance laws many times, usually upwards but sometimes downwards. Papers on the topic have used a variety of combinations of these restrictions as measures of compulsory schooling. We use required years of schooling, defined as the difference between the minimum dropout age and the maximum enrollment age following Lleras-Muney and Goldin and Katz. We follow Acemoglu and Angrist (2001) and Lochner and Moretti (2004) in assigning compulsory attendance laws to people

on the basis of state of birth and the year when the individual was 14 years old (with the exception that the enrollment age is assigned based on the laws in place when the individual was 7 years old). Also, we follow them in creating four indicator variables, depending on whether years of compulsory schooling are 8 or less, 9, 10, and 11 or more.

Our sample is composed of men and women born between 1925 and 1944. We choose this group of cohorts for two reasons. First, many of the changes in compulsory schooling laws were enacted between 1939 and 1958 and so had a major impact on this group. Secondly, the question on age at first marriage is not asked in the Census prior to 1960 or after 1980 so we are limited in terms of which cohorts we can study.

The empirical model can be written as

$$\log(Y_j^*) = X_j^{*\prime} \theta^* + \varepsilon_j, \quad j = 1, \dots, n, \quad (6.1)$$

where Y_j^* denotes age at first marriage for the j^{th} individual in the sample, X_j^* is a vector of explanatory variables including a constant, the compulsory schooling law variables, year of birth dummies, state dummies, and a race dummy, and ε_j an unobserved error term that is uncorrelated with the regressors. There are 3 included compulsory schooling law variables describing the level of compulsory schooling: CA9 (9 years), CA10 (10 years), and CA11 (11 or 12 years). The omitted category is 8 years or less.

There are a few points to note about (6.1): First, it contains fixed cohort effects and state effects. The cohort effects are necessary to allow for secular changes in age at first marriage that may be completely unrelated to compulsory schooling laws. The state effects allow for the fact that variation in the timing of the law changes across states may not have been exogenous to the marriage market (for example, states with strict compulsory schooling laws may be states where people tend to marry late in any case).

The major problem in running this regression is that Y^* is censored for *younger* individuals because we observe age at first marriage only for ever *married* individuals; i.e., for each person we only observe

$$Y_j = \begin{cases} Y_j^* & \text{if } M_j = 1 \\ C_j & \text{if } M_j = 0, \end{cases} \quad (6.2)$$

where C_j is the chronological age and M_j an indicator for ever being married.

There are two elements of the censoring problem: (i) people who do get married at some point in their life but who have never been married at the time of interview, and (ii) people who never get married. Our goal is to address the first problem.¹⁴ The usual approach to dealing with (i) is to restrict the sample to older men and women (e.g., Bergstrom and Schoeni (1996) restrict the sample to persons aged 40–60). This is obviously not a satisfactory solution because it replaces the censoring problem with a truncation problem. In contrast, our approach is to use both young and old persons, acknowledging that age at first marriage is significantly censored for younger women and men. As

¹⁴We cannot solve the second problem as, by definition, it is impossible to construct a refreshment sample for the group that will never marry.

discussed above we use the 1925–1944 cohorts, and these people are aged 16–35 in 1960, and 26–45 in 1970. Clearly, age at first marriage is censored for many of these persons. To deal with this problem, we need a refreshment sample that is not censored and is from the same population as our master sample (aged 16–35 in 1960 and 26–45 in 1970). We obtain this by using individuals from the same cohort: A 16 year old woman in 1960 is considered to be from the same population as a 26 year old woman in 1970, and a 36 year old in 1980. Hence, for women who were between 16–35 in 1960 and 26–45 in 1970, the refreshment sample consists of women aged 36–55 in 1980.

For the group of people aged 36–55 in 1980 to be a suitable refreshment sample, it must possess two characteristics. First, it must be a draw from the same population as the master sample. We consider this to be a reasonable assumption in this case because: (a) they are from the exact same birth cohorts as persons in the master sample; (b) we only use individuals born in the U.S. so immigration is not a problem; (c) we do not include individuals aged more than 55 (and these cohorts were not involved in World War 2 or Vietnam) so mortality is not a major consideration. We report descriptive statistics for our sample in tables 1 and 2 for women and men, respectively. Note that the percentage white, average year-of-birth, and the proportions affected by each compulsory schooling law regime are very similar across Census samples. This is as we would expect given we are tracking a population as they age. On the other hand, the average values of age at first marriage and education differ greatly by Census due to censoring. To further corroborate that we are following samples from the same population, in figure 1 we also present QQ plots for age at first marriage of men and women aged at least 26 that were married before they were 26 years old¹⁵. The linearity of the plots is strong evidence that the uncensored observations in these samples indeed come from the same population.

The second characteristic of a refreshment sample is that it should not have a censoring problem. We examine this issue in table 3. In this table, we track each birth cohort over time, and list the percentage who have never been married. For women, we see that the proportion never married flattens out as women reach their early 30's and it appears that very few women marry for the first time after age 35. Thus, it appears that the refreshment sample for women is approximately free of censoring bias. Men tend to marry at later ages and so there does appear to be some censoring in the refreshment sample for men. However, it impacts a very small proportion of cases; it appears that about 6% of men never marry, and very few cohorts in the refreshment sample have more than 6% of censored observations in 1980. Despite the evidence that there may be some censoring in the 1980 sample, in estimation we treat it as a refreshment sample that has no censored observations.

As mentioned above, we cannot address the second type of censoring (people who never get married) using a refreshment sample approach. Instead, we have taken a few different ad hoc approaches and verify that our results are not very sensitive to the exact method used. The approaches

¹⁵All individuals are aged at least 26 in the 1970 and 1980 samples. To compare 1960 to 1980, we restricted the sample to the oldest 10 cohorts i.e. persons aged at least 26 in 1960 and at least 46 in 1980. This trades off the number of cohorts included against the number of uncensored marriage ages observed.

we have tried are (i) impute age at first marriage as equal to current age for never married individuals in the refreshment sample, and (ii) impute age at marriage for all cases where individuals are not married by 35 (we have tried imputing the age to 55 and 65). We have found that our GMM estimates are reasonably robust to the imputation method used and so we report the results using method (i).

We report the following GMM estimates (which are identical to the EL estimates since here θ^* is just identified) of the coefficients of the compulsory schooling variables and the white dummy in table 4: GMM60, obtained by matching the 1960 master sample with the 1980 refreshment sample to create the enriched dataset, and GMM70, the GMM estimator when the 1970 and 1980 samples are matched. Estimates for men and women are reported separately. Following the procedure described in Section 4.1, see Example 4.2 for an illustration, both estimators were implemented in the GAUSS programming language. Since the consistency of our estimators does not depend upon the extent to which the data are censored, we also expect GMM60 and GMM70 to give similar estimates in finite samples even though censoring is less of a problem in 1970. This is borne out by the evidence summarized in table 4.

An enriched dataset has to, by definition, contain some observations that are not subject to the censoring mechanism. Since age at first marriage is censored from above by chronological age in this application, an enriched dataset here must contain some observations for which age at first marriage is *greater* than chronological age; i.e., loosely speaking, we must have some counterfactual observations for whom we can “look into the future” at the time of interview and see when they first get married. To construct such an enriched dataset by matching, say, the 1960 and 1980 samples, we first create a new variable $\tilde{C}_j = C_j \mathbb{I}(j \in 1960) + (C_j - 20) \mathbb{I}(j \in 1980)$; i.e., \tilde{C}_j is equal to the chronological age for an observation from the 1960 sample, and chronological age less 20 years, otherwise. The enriched observations used to construct GMM60 are then obtained by replacing C_j in (6.2) with \tilde{C}_j . GMM70 is obtained in a similar manner by matching the 1970 and 1980 datasets.

To contrast our GMM estimators with some competing estimators, we also report OLS60, OLS70, TOBIT60, and TOBIT70, the OLS and tobit estimates for each year. Another estimator we consider is OLS80, obtained by doing least squares on just the 1980 sample. It is consistent because the refreshment sample is not censored. Therefore, GMM70 and OLS80 both serve as consistency checks for GMM60. Incidentally, note that although age at first marriage is a continuously distributed random variable, in the data it is recorded in discrete units (years). Therefore, we cannot do censored quantile regression in this application.

First, consider the compulsory schooling estimates for women. The GMM estimates for both 1960 and 1970 are quite similar and suggest that moving from less than 9 years of compulsory schooling to 9 years increases log age at first marriage by about 0.01, implying age at first marriage increases by approximately 1%. The effects for 10 years of compulsory schooling is about 1.5%, and the effects of 11 or more is about 2%. Not surprisingly, these effects are about the same size as one obtains using just the refreshment sample (the 1980 data) because the refreshment sample does not suffer from censoring bias. Note, however, that the GMM estimates are more precisely estimated than the

OLS estimates from 1980, as GMM is optimally using additional information from the 1960 and 1970 samples. The gain in efficiency is bigger for GMM70 than for GMM60, presumably because the 1970 data has less of a censoring problem and hence is more informative¹⁶. The OLS estimates from 1960 and 1970 show signs of bias due to censoring. In particular, the 1960 estimates indicate very large effects of the compulsory schooling laws on age at first marriage. The final two columns in table 4 report tobit estimates. The tobit estimates of the compulsory schooling laws are typically lower than that of the GMM estimators. Also, there is a substantial difference between the tobit estimates for 1960 and the equivalent estimates for 1970, indicating that tobit is performing poorly in this situation.

The estimate of the white dummy for women is also in table 4. The GMM estimates both indicate that whites tend to marry at younger ages than non-whites – the point estimates imply the difference is about 8–9%. Once again, OLS estimates for 1960 and 1970 are very different, suggesting that censoring bias is serious for these samples. The two tobit estimates are again very different from the GMM estimates.

The compulsory schooling and white estimates for men are also in table 4. They differ from the female results in that the GMM estimates only suggest significant effects of 10 years of required schooling (9 years is marginally significant for GMM70). In contrast, the OLS estimates for 60 and 70 show strong significant effects of all the laws on age of first marriage. As in the female sample, the GMM estimates of the white coefficient imply a difference of about 8-9%. Once again, OLS and tobit estimates for 1980 and 1990 are very different, suggesting that censoring bias is serious.

Cohort and state fixed effects were also included in the specification. The estimated cohort effects show how the conditional mean of $\log(\text{age at first marriage})$ varies by birth cohort. The oldest cohort (persons born in 1925) is the excluded dummy in the regression, so the estimate for this group is normalized to zero. Rather than report the coefficients of the cohort dummies, we plot them for women and men in figures 2 and 3, respectively. Not surprisingly, the cohort effects for OLS60 are radically different from the rest. The cohort effects for the rest of the estimators are quite similar to each other.

In summary, we find positive effects of the compulsory schooling laws on age at first marriage. However, the magnitude of the effects are much smaller than would be inferred from ignoring the censoring problem in the 1960 and 1970 data. In contrast, we find large racial differences that are largely obscured in the censored data. Taken together, these demonstrate the importance in this application of using an approach that takes account of censoring. The similarity of the GMM estimates from 1960 and 1970 to each other and to the OLS estimates from 1980 also demonstrates our theoretical result that the proposed estimators are consistent irrespective of the extent of censoring.

¹⁶The difference in the standard errors between OLS80 and the GMM estimators is not that big in this application. We have experimented with reducing the size of the refreshment sample by taking random 10% and 20% subsamples and found a much bigger gain in precision for the GMM estimators over OLS80 (although the resulting estimates are all much less precise than those reported in table 4).

7. CONCLUDING REMARKS AND SOME TOPICS FOR FUTURE RESEARCH

This paper develops efficient semiparametric inference for models with unconditional moment restrictions when the target population is subject to censoring or truncation. Instead of imposing parametric, independence, symmetry, quantile, or special regressor restrictions on the distributions of the underlying random variables, we solve the identification problem created due to the incompleteness of data by using a supplementary sample of observations that are not subject to censoring or truncation. We show how this refreshment sample can be combined with the original dataset of censored or truncated observations to efficiently correct for the effects of partial observation so that all standard GMM and EL based inference goes through. To illustrate our results in an empirical setting, we show how to estimate the effect of changes in compulsory schooling laws on age at first marriage, a variable that is censored for younger individuals, and also demonstrate how refreshment samples in this application can be created by matching cohort information across census datasets.

Of course, the methods developed in this paper are readily applicable in many other applied contexts¹⁷. For example, an important potential application is to the estimation of unemployment durations and re-employment wages subsequent to job displacement. U.S. analyses of the consequences of job displacement have predominantly relied on the Displaced Worker Supplement (DWS) to the Current Population Survey (CPS). However, serious problems arise because many individuals have not become re-employed by the time of the CPS survey so that unemployment durations are censored and re-employment wages are truncated. By using panel data sets such as the Panel Study of Income Dynamics (PSID), one can augment the CPS with a sample that does not have these censoring problems (as individuals are followed for years after displacement) and consistently estimate parameters of interest. We intend to examine this application in future research. The theory developed here can be extended to handle binary response, ordered response, and models involving interval censored or missing data as well. Research on all these topics is also in progress and will be presented in subsequent papers.

APPENDIX A. PROOFS

We only prove the results in Section 4. Results for truncated models in Section 5 can be shown in a similar manner. Henceforth, for notational convenience, let $\hat{\beta} = (\hat{\theta}, \hat{K})_{(p+1) \times 1}$ and $\beta^* = (\theta^*, K_0)$.

Proof of Theorem 4.1. From standard GMM theory we know that $n^{1/2}(\hat{\beta}_{gmm} - \beta^*)$ is asymptotically normal with mean zero and variance $(D'_{f_e} V_{f_e}^{-1} D_{f_e})^{-1}$, where $D_{f_e} = \mathbb{E}_{f_e} \left\{ \frac{\partial \rho(Z, \beta^*)}{\partial \beta} \right\}$ and

¹⁷Applications where refreshment samples are relatively straightforward to construct seem to be those where censoring or truncation can in some sense be regarded as nuisance processes, i.e., where the underlying economic outcomes are not restricted but their measured or recorded versions are. In contrast, it seems hard, at least to us, to non-experimentally construct refreshment samples by matching datasets in applications where censoring or truncation are thought of as being behavioral in origin, i.e., where there are fundamental constraints that bind economic behavior such as those in models of female labor supply or household demand for durable goods.

$V_{f_e} = \mathbb{E}_{f_e} \{\rho(Z, \beta^*) \rho'(Z, \beta^*)\}$. But $V_{f_e} = \begin{bmatrix} V & \gamma \\ \gamma' & \delta \end{bmatrix}$ and, using (4.1), it is easily seen that

$$D_{f_e} = \begin{bmatrix} D & \mathbb{E}_{f^*} \{g(Z^*, \theta^*) \mathbb{I}(Z^* \stackrel{elt}{<} c)\} / K_0 \\ 0'_{p \times 1} & -[1 - F^*(c)] / K_0 \end{bmatrix}.$$

Next, it is straightforward to verify that $\gamma = -\mathbb{E}_{f^*} \{g(Z^*, \theta^*) \mathbb{I}(Z^* \stackrel{elt}{<} c)\} (1 - K_0) / K_0$ and that $\delta = [1 - F^*(c)] (1 - K_0) / K_0$. Using these results, D_{f_e} can be written in a simplified form as

$$D_{f_e} = \begin{bmatrix} D & -\gamma / (1 - K_0) \\ 0'_{p \times 1} & -\delta / (1 - K_0) \end{bmatrix}.$$

Finally, using the partitioned inverse and Sherman-Morrison formulae, a little algebra reveals that

$$D'_{f_e} V_{f_e}^{-1} D_{f_e} = \begin{bmatrix} D' \Omega^{-1} D & 0_{p \times 1} \\ 0'_{p \times 1} & \delta / (1 - K_0)^2 \end{bmatrix}.$$

The desired result follows. \square

Proof of Theorem 4.2. From Qin and Lawless (1994, Theorem 1) we know that $n^{1/2} \{\hat{F}(\xi) - F_e(\xi)\}$ is asymptotically normal with mean zero and variance

$$\text{var}_{f_e} \{\mathbb{I}(Z \leq \xi)\} - \mathbb{E}_{f_e} \{\rho'(Z, \beta^*) \mathbb{I}(Z \leq \xi)\} M_{f_e} \mathbb{E}_{f_e} \{\rho(Z, \beta^*) \mathbb{I}(Z \leq \xi)\},$$

where $M_{f_e} = V_{f_e}^{-1} - V_{f_e}^{-1} D_{f_e} (D'_{f_e} V_{f_e}^{-1} D_{f_e})^{-1} D'_{f_e} V_{f_e}^{-1}$. Using the expressions for D_{f_e} and $D'_{f_e} V_{f_e}^{-1} D_{f_e}$ given in the proof of Theorem 4.1, some tedious but straightforward calculations show that

$$M_{f_e} = \begin{bmatrix} M & -M\gamma/\delta \\ -\gamma'M/\delta & \gamma'M\gamma/\delta^2 \end{bmatrix},$$

where $M = \Omega^{-1} - \Omega^{-1} D (D' \Omega^{-1} D)^{-1} D' \Omega^{-1}$. Therefore,

$$\mathbb{E}_{f_e} \{\rho'(Z, \beta^*) \mathbb{I}(Z \leq \xi)\} M_{f_e} \mathbb{E}_{f_e} \{\rho(Z, \beta^*) \mathbb{I}(Z \leq \xi)\} = \mathbb{E}_{f_e} \{\tilde{g}'(Z) \mathbb{I}(Z \leq \xi)\} M \mathbb{E}_{f_e} \{\tilde{g}(Z) \mathbb{I}(Z \leq \xi)\}.$$

The desired result follows. \square

Proof of Theorem 4.3. First, recall that $\hat{F}^*(\xi) = \sum_{j=1}^n \hat{p}_j(\hat{\beta}_{el}) \mathbb{I}(Z_j \stackrel{elt}{\leq} \xi) \mathbb{I}(Z_j \stackrel{elt}{\neq} c) / a(Z_j, \hat{K}_{el})$. Then, since $1/a(Z_j, \hat{K}_{el}) = \mathbb{I}(Z_j \stackrel{elt}{<} c) + \mathbb{I}(Z_j \stackrel{elt}{<} c) \sim / \hat{K}_{el}$, we can write

$$\hat{F}^*(\xi) = \sum_{j=1}^n \hat{p}_j(\hat{\beta}_{el}) \frac{\mathbb{I}(Z_j \stackrel{elt}{\leq} \xi) \mathbb{I}(Z_j \stackrel{elt}{\neq} c)}{a(Z_j, K_0)} - \frac{(\hat{K}_{el} - K_0)}{\hat{K}_{el} K_0} \sum_{j=1}^n \hat{p}_j(\hat{\beta}_{el}) \mathbb{I}(Z_j \stackrel{elt}{\leq} \xi) \mathbb{I}(Z_j \stackrel{elt}{\neq} c) \mathbb{I}(Z_j \stackrel{elt}{<} c) \sim. \quad (\text{A.1})$$

Next, from (4.5) we know that

$$\hat{K}_{el} = \frac{\sum_{j=1}^n \hat{p}_j(\hat{\beta}_{el}) \mathbb{I}(Z_j \stackrel{elt}{\neq} c) \mathbb{I}(Z_j \stackrel{elt}{<} c) \sim}{\sum_{j=1}^n \hat{p}_j(\hat{\beta}_{el}) \mathbb{I}(Z_j \stackrel{elt}{<} c) \sim}, \quad (\text{A.2})$$

and, by a uniform weak law of large numbers,

$$\begin{aligned} & \sum_{j=1}^n \hat{p}_j(\hat{\beta}_{el}) \mathbb{I}(Z_j \stackrel{elt}{<} c) \sim \xrightarrow{p} \mathbb{E}_{f_e} \{ \mathbb{I}(Z \stackrel{elt}{<} c) \sim \} \\ & \sum_{j=1}^n \hat{p}_j(\hat{\beta}_{el}) \mathbb{I}(Z_j \stackrel{elt}{\neq} c) \mathbb{I}(Z_j \stackrel{elt}{<} c) \sim \xrightarrow{p} \mathbb{E}_{f_e} \{ \mathbb{I}(Z \stackrel{elt}{\neq} c) \mathbb{I}(Z \stackrel{elt}{<} c) \sim \} \\ & \sum_{j=1}^n \hat{p}_j(\hat{\beta}_{el}) \mathbb{I}(Z_j \leq \xi) \mathbb{I}(Z_j \stackrel{elt}{\neq} c) \mathbb{I}(Z_j \stackrel{elt}{<} c) \sim \xrightarrow{p} \mathbb{E}_{f_e} \{ \mathbb{I}(Z \leq \xi) \mathbb{I}(Z \stackrel{elt}{\neq} c) \mathbb{I}(Z \stackrel{elt}{<} c) \sim \}. \end{aligned} \quad (\text{A.3})$$

Hence, since $|\hat{K}_{el} - K_0| = O_p(n^{-1/2})$,

$$\frac{(\hat{K}_{el} - K_0)}{\hat{K}_{el} K_0} \sum_{j=1}^n \hat{p}_j(\hat{\beta}_{el}) \mathbb{I}(Z_j \leq \xi) \mathbb{I}(Z_j \stackrel{elt}{\neq} c) \mathbb{I}(Z_j \stackrel{elt}{<} c) \sim = b(\xi) \mathbb{E}_{f_e} \{ \mathbb{I}(Z \stackrel{elt}{<} c) \sim \} \frac{(\hat{K}_{el} - K_0)}{K_0} + o_p(n^{-1/2}).$$

Therefore, we can rewrite (A.1) as

$$\hat{F}^*(\xi) = \sum_{j=1}^n \hat{p}_j(\hat{\beta}_{el}) \frac{\mathbb{I}(Z_j \leq \xi) \mathbb{I}(Z_j \stackrel{elt}{\neq} c)}{a(Z_j, K_0)} - b(\xi) \mathbb{E}_{f_e} \{ \mathbb{I}(Z \stackrel{elt}{<} c) \sim \} \frac{(\hat{K}_{el} - K_0)}{K_0} + o_p(n^{-1/2}). \quad (\text{A.4})$$

Now, using (A.2) and (4.4), a little algebra reveals that $\mathbb{E}_{f_e} \{ \mathbb{I}(Z \stackrel{elt}{<} c) \sim \} (\hat{K}_{el} - K_0)$ is equal to

$$\frac{\sum_{j=1}^n \hat{p}_j(\hat{\beta}_{el}) [\mathbb{I}(Z_j \stackrel{elt}{\neq} c) \mathbb{I}(Z_j \stackrel{elt}{<} c) \sim \mathbb{E}_{f_e} \{ \mathbb{I}(Z \stackrel{elt}{<} c) \sim \} - \mathbb{I}(Z_j \stackrel{elt}{<} c) \sim \mathbb{E}_{f_e} \{ \mathbb{I}(Z \stackrel{elt}{\neq} c) \mathbb{I}(Z \stackrel{elt}{<} c) \sim \}]}{\sum_{j=1}^n \hat{p}_j(\hat{\beta}_{el}) \mathbb{I}(Z_j \stackrel{elt}{<} c) \sim}. \quad (\text{A.5})$$

Since the term inside the square brackets has mean zero under f_e , as in Qin and Lawless (1994, Theorem 1) we can show that the numerator of (A.5) is $O_p(n^{-1/2})$. Hence, by (A.3),

$$\mathbb{E}_{f_e} \{ \mathbb{I}(Z \stackrel{elt}{<} c) \sim \} (\hat{K}_{el} - K_0) = \sum_{j=1}^n \hat{p}_j(\hat{\beta}_{el}) [\mathbb{I}(Z_j \stackrel{elt}{\neq} c) \mathbb{I}(Z_j \stackrel{elt}{<} c) \sim - K_0 \mathbb{I}(Z_j \stackrel{elt}{<} c) \sim] + o_p(n^{-1/2}). \quad (\text{A.6})$$

But using the fact that $\mathbb{I}(Z_j \stackrel{elt}{\neq} c) \mathbb{I}(Z_j \stackrel{elt}{<} c) = \mathbb{I}(Z_j \stackrel{elt}{<} c)$, it is easily seen that

$$\mathbb{I}(Z_j \stackrel{elt}{\neq} c) \mathbb{I}(Z_j \stackrel{elt}{<} c) \sim - K_0 \mathbb{I}(Z_j \stackrel{elt}{<} c) \sim = K_0 \left\{ \frac{\mathbb{I}(Z_j \stackrel{elt}{\neq} c)}{a(Z_j, K_0)} - 1 \right\}. \quad (\text{A.7})$$

Thus, by (A.4), (A.6), (A.7), and the fact that $\mathbb{E}_{f_e} \{ m(Z, \xi) / a(Z, K_0) \} = F^*(\xi)$, we finally have

$$n^{1/2} \{ \hat{F}^*(\xi) - F^*(\xi) \} = n^{1/2} \sum_{j=1}^n \hat{p}_j(\hat{\beta}_{el}) \left[\frac{m(Z_j, \xi)}{a(Z_j, K_0)} - \mathbb{E}_{f_e} \left\{ \frac{m(Z, \xi)}{a(Z, K_0)} \right\} \right] + o_p(1).$$

Therefore, recalling that $n^{1/2} \{ \hat{F}(\xi) - F_e(\xi) \} = n^{1/2} \sum_{j=1}^n \hat{p}_j(\hat{\beta}_{el}) [\mathbb{I}(Z_j \leq \xi) - \mathbb{E}_{f_e} \{ \mathbb{I}(Z \leq \xi) \}]$, the asymptotic distribution of $n^{1/2} \{ \hat{F}^*(\xi) - F^*(\xi) \}$ can be obtained by replacing $\mathbb{I}(Z \leq \xi)$ in the proof of Theorem 4.2 with $m(Z, \xi) / a(Z, K_0)$. The desired result follows. \square

APPENDIX B. TABLES AND FIGURES

TABLE 1. Descriptive statistics for women by year

	Mean	Std. Dev.	Min	Max
<u>1960 (220730 observations)</u>				
Birth Cohort	1934.62	5.98	1925	1944
Age	25.38	5.98	16	35
Age at First Marriage	19.68	3.59	14	35
Never Married	0.29	0.45	0	1
White	0.88	0.32	0	1
≤ 8 Years of Schooling Required	0.19	0.39	0	1
9 Years of Schooling Required	0.66	0.47	0	1
10 Years of Schooling Required	0.08	0.27	0	1
≥ 11 Years of Schooling Required	0.07	0.26	0	1
<u>1970 (216036 observations)</u>				
Birth Cohort	1934.69	5.94	1925	1944
Age	35.31	5.94	26	45
Age at First Marriage	21.23	5.19	14	45
Never Married	0.07	0.25	0	1
White	0.88	0.32	0	1
≤ 8 Years of Schooling Required	0.19	0.39	0	1
9 Years of Schooling Required	0.66	0.47	0	1
10 Years of Schooling Required	0.08	0.27	0	1
≥ 11 Years of Schooling Required	0.07	0.26	0	1
<u>1980 (223903 observations)</u>				
Birth Cohort	1934.73	5.95	1925	1944
Age	45.28	5.95	36	55
Age at First Marriage	22.07	7.01	12	55
Never Married	0.05	0.22	0	1
White	0.88	0.33	0	1
≤ 8 Years of Schooling Required	0.19	0.39	0	1
9 Years of Schooling Required	0.66	0.47	0	1
10 Years of Schooling Required	0.08	0.27	0	1
≥ 11 Years of Schooling Required	0.07	0.26	0	1

TABLE 2. Descriptive statistics for men by year

	Mean	Std. Dev.	Min	Max
<u>1960 (213184 observations)</u>				
Birth Cohort	1934.69	6.00	1925	1944
Age	25.31	6.00	16	35
Age at First Marriage	21.36	3.95	14	35
Never Married	0.42	0.49	0	1
White	0.89	0.31	0	1
≤ 8 Years of Schooling Required	0.19	0.39	0	1
9 Years of Schooling Required	0.66	0.47	0	1
10 Years of Schooling Required	0.08	0.27	0	1
≥ 11 Years of Schooling Required	0.07	0.26	0	1
<u>1970 (207129 observations)</u>				
Birth Cohort	1934.71	5.94	1925	1944
Age	35.29	5.94	26	45
Age at First Marriage	23.82	5.26	14	45
Never Married	0.10	0.30	0	1
White	0.90	0.30	0	1
≤ 8 Years of Schooling Required	0.19	0.39	0	1
9 Years of Schooling Required	0.66	0.47	0	1
10 Years of Schooling Required	0.08	0.27	0	1
≥ 11 Years of Schooling Required	0.07	0.26	0	1
<u>1980 (212244 observations)</u>				
Birth Cohort	1934.80	5.93	1925	1944
Age	45.20	5.93	36	55
Age at First Marriage	24.95	7.26	12	55
Never Married	0.07	0.25	0	1
White	0.89	0.31	0	1
≤ 8 Years of Schooling Required	0.19	0.39	0	1
9 Years of Schooling Required	0.66	0.47	0	1
10 Years of Schooling Required	0.08	0.27	0	1
≥ 11 Years of Schooling Required	0.07	0.26	0	1

TABLE 3. Proportion censored by cohort and year

age in 1960	<u>% of women censored</u>			<u>% of men censored</u>		
	1960	1970	1980	1960	1970	1980
16	94	13	7	99	20	9
17	88	11	6	98	17	8
18	75	10	6	95	15	8
19	59	9	6	87	13	8
20	46	8	6	75	12	7
21	34	7	5	62	10	7
22	25	7	5	50	10	6
23	19	7	5	40	9	6
24	15	6	5	32	8	7
25	13	6	5	27	9	6
26	11	5	5	22	8	6
27	9	6	4	19	7	6
28	9	5	5	16	7	5
29	9	5	5	15	7	6
30	8	5	4	13	7	6
31	7	5	4	12	7	6
32	6	5	4	11	7	6
33	6	5	5	11	7	6
34	6	5	4	10	7	6
35	6	5	5	9	6	6
36	6	5	4	8	6	6
37	6	6	5	8	6	6
38	5	5	5	8	6	6
39	6	5	5	8	6	6
40	6	6	5	7	6	6

TABLE 4. Effects of compulsory schooling laws and race on log(age at first marriage). Also included in the specification, but not reported in this table, are a constant, year-of-birth indicators, and state dummies.

Women	OLS60	OLS70	OLS80	GMM60	GMM70	TOBIT60	TOBIT70
9 Years Schooling Req'd.	.0157* (.0016)	.0080* (.0022)	.0096* (.0025)	.0102* (.0024)	.0094* (.0022)	.0029 (.0018)	.0077* (.0021)
10 Years Schooling Req'd.	.0232* (.0021)	.0112* (.0030)	.0146* (.0035)	.0150* (.0034)	.0129* (.0031)	.0075* (.0026)	.0103* (.0031)
11+ Years Schooling Req'd.	.0456* (.0038)	.0317* (.0052)	.0184* (.0061)	.0188* (.0060)	.0223* (.0053)	.0157* (.0049)	.0299* (.0057)
White	-.0261* (.0012)	-.0476* (.0018)	-.0827* (.0021)	-.0927* (.0020)	-.0808* (.0019)	-.0393* (.0014)	-.0534* (.0016)
Men	OLS60	OLS70	OLS80	GMM60	GMM70	TOBIT60	TOBIT70
9 Years Schooling Req'd.	.0114* (.0015)	.0073* (.0022)	.0031 (.0026)	.0046 (.0024)	.0061* (.0023)	-.0028 (.0019)	.0062* (.0022)
10 Years Schooling Req'd.	.0205* (.0019)	.0120* (.0029)	.0130* (.0035)	.0131* (.0033)	.0137* (.0031)	.0052 (.0029)	.0109* (.0031)
11+ Years Schooling Req'd.	.0359* (.0035)	.0152* (.0053)	.0049 (.0063)	.0028 (.0060)	.0070 (.0056)	.0055 (.0053)	.0121* (.0057)
White	-.0156* (.0011)	-.0444* (.0018)	-.0792* (.0021)	-.0826* (.0020)	-.0792* (.0019)	-.0301* (.0016)	-.0515* (.0017)

Standard errors in parenthesis. A “*” superscript denotes that effect is significant at 5% level.




FIGURE 1. QQ plots of age at first marriage for individuals aged at least 26 that are uncensored; i.e., those who married before age 26.




FIGURE 2. Cohort effects for women.




FIGURE 3. Cohort effects for men.

REFERENCES

- ACEMOGLU, D., AND J. ANGRIST (2001): "How large are human capital externalities? Evidence from compulsory schooling laws," in *NBER Macroeconomics Annual 2000*, ed. by B. S. Bernanke, and K. Rogoff. MIT Press.
- AKERLOF, G. A. (1998): "Men without children," *Economic Journal*, 108, 287–309.
- AMEMIYA, T. (1985): *Advanced econometrics*. Harvard University Press.
- BERGSTROM, T., AND R. F. SCHOENI (1996): "Income prospects and age-at-marriage," *Journal of Population Economics*, 9, 115–130.
- BHATTACHARYA, P., H. CHERNOFF, AND S. YANG (1983): "Nonparametric estimation of the slope of a truncated regression," *The Annals of Statistics*, 11, 505–514.
- BLUNDELL, R. W., AND R. J. SMITH (1993): "Simultaneous microeconomic models with censored or qualitative dependent variables," in *Econometrics*. Amsterdam: North-Holland, vol. 11 of *Handbook of Statistics*, pp. 117–143.
- BROWN, B. W., AND W. K. NEWEY (2002): "GMM, efficient bootstrapping, and improved inference," *Journal of Business and Economic Statistics*, 20, 507–517.
- BUCHINSKY, M., AND J. HAHN (1998): "An alternative estimator for the censored quantile regression model," *Econometrica*, 66, 653–672.
- CARROLL, R., D. RUPPERT, AND L. STEFANSKI (1995): *Measurement error in nonlinear models*. Chapman & Hall/CRC.
- CHAMBERLAIN, G. (1986): "Asymptotic efficiency in semiparametric models with censoring," *Journal of Econometrics*, 32, 189–218.
- (1987): "Asymptotic efficiency in estimation with conditional moment restrictions," *Journal of Econometrics*, 34, 305–334.
- CHEN, S., AND S. KHAN (2001): "Estimation of a partially linear censored regression model," *Econometric Theory*, 17, 567–590.
- CHEN, X., H. HONG, AND E. TAMER (2004): "Measurement error models with auxiliary data," *Review of Economic Studies*, Forthcoming.
- COALE, A. (1971): "Age patterns of marriage," *Population Studies*, 25, 193–214.
- COSSLETT, S. (1981): "Efficient estimation of discrete choice models," in *Structural analysis of discrete data with econometric applications*, ed. by C. F. Manski, and D. McFadden. Cambridge, MA: MIT Press, pp. 51–111.
- (1987): "Efficiency bounds for distribution free estimators of binary choice and censored regression models," *Econometrica*, 55, 559–585.
- (1991): "Distribution-free estimator of a regression model with sample selectivity," in *Nonparametric and semiparametric methods in econometrics and statistics. Proceedings of the fifth international symposium in Economic Theory and Econometrics*, ed. by W. A. Barnett, J. L. Powell, and G. Tauchen, pp. 175–197. Cambridge University Press.
- DEMPSTER, A., N. LAIRD, AND D. RUBIN (1977): "Maximum likelihood from incomplete data via the EM algorithm," *Journal of the Royal Statistical Society, Series B*, 39, 1–38.
- DUNCAN, G. (1986): "A semiparametric censored regression estimator," *Journal of Econometrics*, 32, 5–34.
- FERNANDEZ, L. (1986): "Nonparametric maximum likelihood estimation of censored regression models," *Journal of Econometrics*, 32, 35–57.
- GOLDIN, C., AND L. F. KATZ (2002): "The power of the Pill: Oral contraceptives and womens career and marriage decisions," *Journal of Political Economy*, 110, 730–770.
- (2003): "Mass secondary schooling and the State: The role of state compulsion in the high school movement," NBER Working Paper 10075.
- HANSEN, L. P. (1982): "Large sample properties of generalized methods of moments estimators," *Econometrica*, 50, 1029–1054.

- HAUSMAN, J. A., AND D. A. WISE (1976): "The evaluation of results from truncated samples: The New Jersey negative income tax experiment," *Annals of Economic and Social Measurement*, 5, 421–445.
- (1977): "Social experimentation, truncated distributions, and efficient estimation," *Econometrica*, 45, 919–938.
- HECKMAN, J. J. (1976): "The common structure of statistical models of truncation, sample selection and limited dependent variables and a simple estimator for such models," *Annals of Economic and Social Measurement*, 5/4, 475–492.
- (1979): "Sample selection bias as a specification error," *Econometrica*, 47, 153–161.
- HELLERSTEIN, J., AND G. W. IMBENS (1999): "Imposing moment restrictions from auxiliary data by weighting," *Review of Economics and Statistics*, 81, 1–14.
- HIRANO, K., G. W. IMBENS, G. RIDDER, AND D. B. RUBIN (2001): "Combining panel data sets with attrition and refreshment samples," *Econometrica*, 69, 1645–1659.
- HONG, H., AND E. TAMER (2003): "Inference in censored models with endogenous regressors," *Econometrica*, 71, 905–932.
- HONORÉ, B., AND J. L. POWELL (1994): "Pairwise difference estimators of censored and truncated regression models," *Journal of Econometrics*, 64, 241–278.
- HOROWITZ, J. L. (1986): "A distribution-free least squares estimator for censored linear regression models," *Journal of Econometrics*, 32, 59–84.
- (1988): "Semiparametric M-estimation of censored linear regression models," *Advances in Econometrics*, 7, 45–83.
- HOROWITZ, J. L., AND C. F. MANSKI (1998): "Censoring of outcomes and regressors due to survey nonresponse: Identification and estimation using weights and imputations," *Journal of Econometrics*, 84, 37–58.
- IMBENS, G. W. (1997): "One-step estimators for over-identified generalized method of moments models," *Review of Economic Studies*, 64, 359–383.
- IMBENS, G. W., AND T. LANCASTER (1994): "Combining micro and macro data in microeconomic models," *Review of Economic Studies*, 61, 655–680.
- KHAN, S., AND A. LEWBEL (2003): "Weighted and two stage least squares estimation of semiparametric truncated regression models," Manuscript.
- KHAN, S., AND J. L. POWELL (2001): "Two step estimation of semiparametric censored regression models," *Journal of Econometrics*, 103, 73–110.
- KITAMURA, Y. (1997): "Empirical likelihood methods with weakly dependent processes," *Annals of Statistics*, 25, 2084–2102.
- (2001): "Asymptotic optimality of empirical likelihood for testing moment restrictions," *Econometrica*, 69, 1661–1672.
- KOBALL, H. (1998): "Have African American men become less committed to marriage? Explaining the twentieth century racial cross-over in mens marriage timing," *Demography*, 35, 251–258.
- KORENMAN, S., AND D. NEUMARK (1991): "Does marriage really make men more productive," *Journal of Human Resources*, 26, 282–307.
- (1992): "Marriage, motherhood, and wages," *Journal of Human Resources*, 27, 233–255.
- LANCASTER, T. (1992): "A paradox in choice based sampling," Manuscript.
- LEE, M. (1993a): "Quadratic mode regression," *Journal of Econometrics*, 57, 1–19.
- (1993b): "Winsorized mean estimator for censored regression model," *Econometric Theory*, 8, 368–382.
- LLERAS-MUNEY, A. (2001): "Were compulsory attendance and child labor laws effective? An analysis from 1915 to 1939," *Journal of Law and Economics* (forthcoming).
- (2002): "The relationship between education and adult mortality in the United States," NBER working paper 8986.

- LOCHNER, L., AND E. MORETTI (2004): "The effect of education on crime: Evidence from prison inmates, arrests, and self-reports," *American Economic Review*, 94, 155–189.
- MADDALA, G. (1983): *Limited-dependent and qualitative variables in econometrics*. Cambridge University Press.
- MANSKI, C. F., AND S. R. LERMAN (1977): "The estimation of choice probabilities from choice based samples," *Econometrica*, 45, 1977–1988.
- MOON, C.-G. (1989): "A Monte Carlo comparison of semiparametric tobit estimators," *Journal of Applied Econometrics*, 4, 361–382.
- NAN, B., M. EMOND, AND J. A. WELLNER (2002): "Information bounds for Cox regression models with missing data," Manuscript.
- NAWATA, K. (1990): "Robust estimation based on grouped-adjusted data in censored regression models," *Journal of Econometrics*, 43, 337–362.
- NEVO, A. (2003): "Using weights to adjust for sample selection when auxiliary information is available," *Journal of Business and Economic Statistics*, 21, 43–52.
- NEWKEY, W. K. (1988): "Efficient estimation of Tobit models under symmetry," in *Nonparametric and semiparametric methods in econometrics and statistics. Proceedings of the fifth international symposium in Economic Theory and Econometrics*, ed. by W. A. Barnett, J. Powell, and G. Tauchen, pp. 291–336. Cambridge University Press.
- (2001): "Conditional moment restrictions in censored and truncated regression models," *Econometric Theory*, 17, 863–888.
- NEWKEY, W. K., AND D. MCFADDEN (1994): "Large sample estimation and hypothesis testing," in *Handbook of Econometrics, vol. IV*, ed. by R. Engle, and D. McFadden. Elsevier Science B.V., pp. 2111–2245.
- NEWKEY, W. K., AND J. L. POWELL (1990): "Efficient estimation of linear and type I censored regression models under conditional quantile restrictions," *Econometric Theory*, 6, 295–317.
- NEWKEY, W. K., AND R. J. SMITH (2003): "Higher order properties of GMM and generalized empirical likelihood estimators," *Econometrica*, Forthcoming.
- OWEN, A. (1988): "Empirical likelihood ratio confidence intervals for a single functional," *Biometrika*, 75(2), 237–249.
- (2001): *Empirical likelihood*. Chapman and Hall/CRC.
- POWELL, J. L. (1983): "The asymptotic normality of two stage least absolute deviations estimators," *Econometrica*, 51, 1569–1575.
- (1984): "Least absolute deviations estimation for the censored regression model," *Journal of Econometrics*, 25, 303–325.
- (1986a): "Censored regression quantiles," *Journal of Econometrics*, 32, 143–155.
- (1986b): "Symmetrically trimmed least squares estimation for tobit models," *Econometrica*, 54, 1435–1460.
- (1994): "Estimation of semiparametric models," in *Handbook of Econometrics, vol. IV*, ed. by R. Engle, and D. McFadden. Elsevier Science B.V., pp. 2443–2521.
- QIN, J., AND J. LAWLESS (1994): "Empirical likelihood and general estimating equations," *Annals of Statistics*, 22, 300–325.
- RIDDER, G. (1992): "An empirical evaluation of some models for non-random attrition in panel data," *Structural Change and Economic Dynamics*, 3, 337–355.
- ROBINS, J. M., F. HSIEH, AND W. NEWKEY (1995): "Semiparametric efficient estimation of a conditional density with missing or mismeasured covariates," *Journal of the Royal Statistical Society, Series B*, 57, 409–424.
- ROBINS, J. M., A. ROTNITZKY, AND L. P. ZHAO (1994): "Estimation of regression coefficients when some regressors are not always observed," *Journal of the American Statistical Association*, 89, 846–866.
- SMITH, R. J. (1997): "Alternative semi-parametric likelihood approaches to generalized method of moments estimation," *Economic Journal*, 107, 503–519.

- SMITH, R. J., AND R. W. BLUNDELL (1986): "An exogeneity test for a simultaneous equation tobit model with an application to labor supply," *Econometrica*, 54, 679–685.
- TITTERINGTON, D. (1983): "Kernel-based density estimation using censored, truncated or grouped data," *Communications in Statistics, Theory and Methods*, 12, 2151–2167.
- TITTERINGTON, D., AND G. MILL (1983): "Kernel-based density estimates from incomplete data," *Journal of the Royal Statistical Society, Series B*, 45, 258–266.
- TRIPATHI, G. (2003): "GMM and empirical likelihood with stratified data," Manuscript. Department of Economics, University of Wisconsin-Madison.
- (2004): "Moment based inference with stratified data," Manuscript. Department of Economics, University of Connecticut-Storrs.
- TSUI, K.-L., N. P. JEWELL, AND C. WU (1988): "A nonparametric approach to the truncated regression problem," *Journal of the American Statistical Association*, 83, 785–792.
- VARDI, Y. (1985): "Empirical distributions in selection biased models," *Annals of Statistics*, 13, 178–203.
- VELLA, F. (1998): "Estimating models with sample selection bias: A survey," *Journal of Human Resources*, 33, 127–172.
- WANSBEEK, T., AND E. MEIJER (2000): *Measurement error and latent variables in econometrics*. North-Holland, Amsterdam.

DEPARTMENT OF ECONOMICS, UNIVERSITY OF CALIFORNIA, LOS ANGELES, CA-90095.

E-mail address: devereux@econ.ucla.edu

URL: www.econ.ucla.edu/devereux

DEPARTMENT OF ECONOMICS, UNIVERSITY OF CONNECTICUT, STORRS, CT-06269.

E-mail address: gautam.tripathi@uconn.edu

URL: web.uconn.edu/tripathi